

PCTWORLD INTELLECTUAL PRO
International

INTERNATIONAL APPLICATION PUBLISHED UNDER

WO 9606944A1

(51) International Patent Classification ⁶ : C12P 19/34, C07H 21/02	A1	(11) International Publication Number: WO 96/06944 (43) International Publication Date: 7 March 1996 (07.03.96)
(21) International Application Number: PCT/US95/10813 (22) International Filing Date: 25 August 1995 (25.08.95) (30) Priority Data: 08/299,498 1 September 1994 (01.09.94) US (71) Applicant: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US). (72) Inventors: SZOSTAK, Jack, W.; 308 Commonwealth Avenue, Boston, MA 02115 (US). LORSCH, Jon, R.; 22 Concord Avenue #4, Cambridge, MA 02138 (US). WILSON, Charles; 180 Mountain Springs Drive, Bonny Doon, CA 95060 (US). (74) Agent: LECH, Karen, F.; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: NOVEL RIBOZYMES AND NOVEL RIBOZYME SELECTION SYSTEMS (57) Abstract The invention concerns a method for creating, identifying, and isolating ribozymes capable of binding a selected ligand and catalyzing a reaction involving the selected ligand. The method entails sequential selections for ligand binding molecules and catalytic molecules. The invention also includes novel ribozymes produced by these methods.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

NOVEL RIBOZYMES AND NOVEL RIBOZYME SELECTION SYSTEMSBackground of the Invention

5 This invention relates to novel ribozyme molecules and methods for their identification and isolation.

 This invention was made with Government support under Contract #R01-GM45315-02 awarded by the National Institutes of Health. The Government has certain rights
10 in this invention.

 Both the genetic and enzymatic components of the earliest cells are thought to have been RNA molecules, because RNA is the only known macromolecule that can both encode information in a heritable form, and act as a
15 biocatalyst (Joyce, Nature 338:217, 1989). It has been proposed that modern metabolism evolved prior to the evolution of encoded protein synthesis, and that early ribozyme-catalyzed metabolic transformations form the basis of our present protein-catalyzed metabolism (Benner
20 et al., Proc. Natl. Acad. Sci. USA 86:7054, 1989). This proposal requires that ribozymes should be able to catalyze a broad range of chemical transformations. However, to date, known natural ribozymes, including the group I and group II introns, RNase P, and the hammerhead
25 and hairpin RNAs, have been shown to catalyze only a restricted range of reactions involving the RNA sugar-phosphate backbone (Wilson and Szostak, Curr. Opin. Struct. Biol. 2:749, 1992).

Summary of the Invention

30 The invention concerns a method for creating, identifying, and isolating catalytic RNA molecules capable of binding a ligand and catalyzing a reaction modifying the catalytic RNA (or other substrate). The method entails sequential selections for ligand binding
35 RNA molecules and catalytic RNA molecules.

- 2 -

The ribozymes isolated by the method of the invention are capable of catalyzing reactions normally catalyzed by enzymes. Previously, the art disclosed ribozymes capable of catalyzing reactions involving the RNA sugar-phosphate backbone, e.g., phosphodiester transfer reactions and hydrolysis of nucleic acids. The methods of the invention can be used to create ribozymes capable of carrying out reactions on the RNA sugar-phosphate backbone. In addition, however, ribozymes created by the method of the invention can catalyze reactions other than hydrolysis and transesterification, thereby increasing the range of systems for which the catalytic ribozymes and the catalytic ribozyme selection systems of the invention are useful.

15 The methods of the invention entail sequential in vitro selections using pools of RNA molecules which include one or more regions of random sequence. Because catalysis of a complex reaction demands both the ability to bind a non-RNA ligand and the preferential

20 stabilization of the transition state configuration of the reactants, the number of functional ribozymes in a pool of RNA having one or more regions of random sequence may be vanishingly small. The methods of the invention overcome this difficulty through the use of sequential

25 selections. The method of the invention entails at least two selections steps: a binding selection step for identifying in a pool of random RNA molecules those RNA molecules which are capable of binding the selected ligand and a catalysis selection step for identifying in

30 a pool of substrate binding RNA molecules (or sequence variants of such RNA molecules) those which are capable of catalyzing a reaction which modifies the catalytic RNA (or other substrate). After each selection step, an amplification step is performed. In this amplification

35 step, the selected molecules are amplified using PCR. Of

- 3 -

course, as explained more fully below, the binding selection step and the catalysis selection step may include one, two, or more rounds of selection and amplification. After each round, the pool of molecules

5 is enriched for those having the desired binding or catalysis activity. Thus, the methods of the invention effectively entail three steps: 1) selection of RNA molecules capable of binding a chosen ligand from a pool of RNA molecules having a region of random sequence; 2)

10 generation of a pool of RNA molecules which have a ligand binding sequence which is based on the identified ligand binding sequence of ligand-binding RNA molecules selected in step 1 as well as a region of random sequence; and 3)

15 selection of RNA molecules exhibiting catalytic activity which modifies the RNA molecule itself or a substrate attached to the catalytic RNA. To identify catalytic RNA molecules one must tag the active molecules so that they may be partitioned from the inactive ones. This tagging is most straightforward when the reaction catalyzed by

20 the RNA molecule modifies the catalytic RNA molecule itself. This modification can involve the formation of a chemical bond, the breaking of a chemical bond, or both. Often the modification attaches one or more new atoms to the RNA. Other desirable modifications remove one or

25 more atoms from the RNA. To be useful for tagging the modification must render the modified molecules distinguishable from non-modified molecules. Tagging can also be accomplished by modification of a substrate attached to the catalytic RNA molecule. If all of the

30 molecules in the pool are attached to a substrate molecule, those RNA molecules which can catalyze a reaction modifying the attached substrate can be partitioned from the RNA molecules which do not carry out the modification.

- 4 -

Of course, one may find that a pool of catalytic RNA molecules is capable of carrying out a number of different modifications.

The selected ligand can include small molecules
5 such as drugs, metabolites, cofactors, toxins, and transition state analogs. Possible ligands also include proteins, polysaccharides, glycoproteins, hormones, receptors, lipids, and natural or synthetic polymers. Preferably, for therapeutic applications, binding of the
10 ligand and catalysis takes place in aqueous solution under physiological or near physiological salt conditions, temperature, and pH.

It is important to note that the ligand used to identify ligand-binding RNA molecules may be, but does
15 not have to be, the same ligand which is used in the catalyst selection step. One may wish, for example, to isolate ligand-binding RNA molecules using a first ligand (e.g., ATP) and then isolate catalytic RNA molecules with a second ligand (e.g., ATP- γ -S) which can bind to the
20 same ligand binding region.

As mentioned above, the method of the invention entails at least two selection steps. In the first step, RNA molecules capable of binding the chosen ligand are selected from a pool of RNA molecules which include one
25 or more regions of random sequence. In the second selection step, RNA molecules capable of catalyzing a reaction modifying the RNA (or other substrate) are chosen from a second pool of random RNA molecules whose sequence is based on the sequence of one or more ligand
30 binding RNAs identified in the first selection step.

"Random RNAs" and "random sequence" are general terms used to describe molecules or sequences which have one or more regions of "fully random sequence" and/or one or more regions of "partially random sequence." Such
35 molecules may also include one or more regions of

- 5 -

"defined sequence." "Fully random sequence" is sequence in which there is a roughly equal probability of each of A, T, C, and G being present at each position in the sequence. Of course, the limitations of some of the methods used to create nucleic acid molecules make it rather difficult to create fully random sequences in which the probability of each nucleotide occurring at each position is absolutely equal. Accordingly, sequences in which the probabilities are roughly equal are considered fully random sequences. In "partially random sequences" and "partially randomized sequences," rather than there being a 25% chance of each of A, T, C, and G being present at each position, there are unequal probabilities. For example, in a partially random sequence, there may be a 70% chance of A being present at a given position and a 10% chance of each of T, C, and G being present. Further, the probabilities can be the same or different at each position within the partially randomized region. Thus, a partially random sequence may include one or more positions at which the sequence is fully random and one or more positions at which the sequence is defined. Such partially random sequences are particularly useful when one wishes to make variants of a known sequence. For example, if one knows that a particular 20 base sequence binds the selected ligand and that positions 2, 3, 4, 12, 13, and 15-20 are critical for binding, one could prepare a partially random version of the 20 base sequence in which the bases at positions 2, 3, 4, 12, 13, and 15-20 are the same as in the known ligand binding sequence and the other positions are fully randomized. Alternatively, one could prepare a partially random sequence in which positions 2, 3, 4, 12, 13, and 15-20 are partially randomized, but with a strong bias towards the bases found at each position in the original molecule, with all of the other positions being fully

- 6 -

randomized. This type of partially random sequence is desirable in pools of molecules from which catalytic RNAs are being selected.

As discussed below, the sequence of any randomized
5 region may be further randomized by mutagenesis during one or more amplification steps as part of a process referred to as in vitro evolution.

It is desirable to have one, preferably two, regions of "defined sequence". Defined sequence is
10 sequence selected or known by the creator of the molecule. Such defined sequence regions are useful for isolating and amplifying the nucleic acid because they are recognized by defined complementary primers. The defined primers can be used to isolate or amplify
15 sequences having the corresponding defined sequences. The defined sequence regions preferably flank the randomized regions. The defined region or regions can also be intermingled with the randomized regions. Both the random and specified regions can be of any desired
20 length.

In the first step, nucleic acids capable of binding the ligand are identified. Beginning with a pool of nucleic acids which include one or more regions of random sequence, the method for isolating ligand-binding
25 molecules includes contacting the pool of nucleic acid with the substrate under conditions which are favorable for binding, partitioning nucleic acids which have bound the substrate from those which have not, dissociating bound nucleic acids and substrate, amplifying the nucleic
30 acids (e.g., using PCR) which were previously bound, and, if desired, repeating the steps of binding, partitioning, dissociating, and amplifying any desired number of times.

Several cycles of selection (binding, partitioning, dissociating, and amplifying) are desirable
35 because after each round the pool is more enriched for

- 7 -

substrate binding nucleic acids. One can perform additional cycles of selection until no substantial improvement in substrate binding is observed. Of course, one can also perform far fewer cycles of selection.

5 In many cases, sequencing of nucleic acids isolated after one or more rounds of partitioning and amplification will reveal the presence of a number of different nucleic acids. One or more of these nucleic acids can be used in the pool of nucleic acids from which
10 catalytic nucleic acids are isolated in the second selection of the method of the invention. Alternatively, the pool for the second phase can be composed of one or more nucleic acids having sequences based on the sequences of the nucleic acids identified in the binding
15 selection. For example, sequencing of the nucleic acids which bind the substrate may suggest one or more regions of consensus sequence, i.e., sequences which appear to be important for binding. The pool of molecules used for selection of catalytic molecules may then include nucleic
20 acids whose sequence is based on this consensus sequence. One may also employ a partially randomized sequence based on the consensus sequence. This may permit the isolation of improved binding domains. It can also permit alterations of the binding domain which may be desirable
25 for improved catalysis. Of course, as discussed above, the degree of randomization of the consensus sequence is generally quite low. The consensus sequence region, randomized or not, may be interspersed with and/or flanked by additional randomized regions. Thus, the
30 sequences of the molecules in the pool of nucleic acids molecules used in the catalysis selection step can differ from that of the molecule(s) identified in the substrate selection step as molecules capable of binding the desired substrate.

- 8 -

Those skilled in the art can readily identify ligand-binding consensus sequences by sequencing a number of ligand-binding RNA molecules and comparing their sequences. In some cases such sequencing and comparison
5 will reveal the presence of a number of different classes of ligand binding sequences (aptamers). In these circumstances it may be possible to identify a core sequence which is common to most or all classes. This core sequence or variants thereof can be used as the
10 starting point for the catalysis selection. By "variant" of a ligand binding sequence is meant a sequence created by partially randomizing a ligand binding sequence.

The size of the randomized regions employed should be adequate to provide a substrate binding site in the
15 case of the binding selection step. Thus, the randomized region used in the initial selection preferably includes between 15 and 60 nucleotides, more preferably between 20 and 40 nucleotides. The randomized region or regions used for the catalysis selection step should be of
20 sufficient length to provide a reasonable probability of being able to include catalytic activity.

The probability that any given RNA sequence of 30, 50, 100, or even 400 bases includes a region capable of binding a chosen substrate is very low. Similarly, the
25 probability that a given RNA sequence which includes a region capable of binding a chosen substrate also has a region capable of catalyzing a reaction involving the chosen substrate is very low. Because of this each, of the two selection steps preferably begins with a pool of
30 molecules which is large enough and random enough to include molecules which can bind the chosen substrate in the case of the binding selection or catalyze a reaction involving the chosen substrate in the case of the catalysis selection. Accordingly, the molecules used in
35 each initial pool include at least one fully random

- 9 -

sequence region. Binding sites may occur at a frequency of 10^{-10} to 10^{-15} in random sequences. Thus, pool sizes are preferably greater than 10^{10} .

It is generally not practical to prepare a
5 population of molecules which includes all of the possible sequences of a particular random sequence. However, even where one has a population of no more than 10^{15} different molecules out of 10^{60} potential sequences, one can isolate molecules having a desired binding or
10 catalytic activity.

The catalysis selection step involves identifying RNAs which catalyze a reaction involving the chosen ligand. The pool of molecules used at the outset of this selection step generally is composed of molecules having
15 one or more defined or partially randomized sequences which are designed to bind to the chosen ligand ("ligand binding region") as well as a second random sequence region, preferably fully randomized which serves as the source of potentially catalytic sequences. The ligand
20 binding region included in the molecules in this catalysis selection pool can have a sequence which is identical to an identified ligand binding sequence identified in the binding selection phase. Alternatively the sequence of this region can be based on the consensus
25 sequence of a number of substrate binding regions identified in the first step. The region may also be a partially randomized sequenced based on either a particular substrate binding sequence or substrate binding consensus sequence. Of course, the molecules
30 also preferably include one or more defined sequence regions which can bind isolation or amplification primers.

In order to identify molecules having catalytic activity there must be a means for partitioning those RNA
35 molecules which have catalyzed a reaction modifying the

- 10 -

RNA molecule (or a substrate attached to the RNA) from those which have not. The selection can be accomplished using affinity columns which will bind modified, but not unmodified molecules. Alternatively, one can employ an
5 antibody which recognizes the modified, but not unmodified molecules. It is also possible to chemically convert modified, but not unmodified ligand, to a compound which will bind selectively to an affinity column or other selective binding material (e.g., an
10 antibody).

In many cases the catalytic RNA will itself be chemically altered (modified) by the reaction it catalyzes. This alteration can then form the basis for selecting catalytic molecules.

15 In many cases it may be possible to alter such catalytic RNA molecules so that instead of being self-modifying they modify a second molecule.

As will be apparent from the examples below there are a number of means for partitioning catalytic
20 molecules from non-catalytic or less catalytic molecules.

It may be desirable to increase the stringency of a selection step in order to isolate more desirable molecules. The stringency of the binding selection step can be increased by decreasing ligand concentration. The
25 stringency of the catalysis selection step can be increased by decreasing the ligand concentration or the reaction time.

One can covalently link a molecule to be modified to RNA so that catalytic RNA molecules can be isolated by
30 isolating the modified molecule. For example, one might wish to find RNAs capable of oxidizing compound A. This might be accomplished by isolating RNA molecules capable of binding a redox co-factor (NAD, FAD, or NADP). A pool of random RNAs is then created which are capable of

- 11 -

binding the cofactor. Compound A is then covalently attached to the RNA molecules in this pool and a selection is carried out which isolates molecules having the oxidized form of compound A. Methods for linking
5 various compounds to RNA are well known to those skilled in the art and include the use of a thiophosphate group and the use of amines linked via a 5' phosphate.

Of course, in some cases a catalytic RNA which is capable of self-modification or modification of an
10 attached substrate may also be able to perform the "trans" reaction. Such trans acting molecules modify an RNA other than themselves or modify the substrate even when it is not attached to the catalytic RNA.

In one aspect, therefore, the invention features a
15 method for producing a catalytic RNA molecule capable of binding a first ligand and catalyzing a chemical reaction modifying the catalytic RNA molecule. The method includes the following steps:

- a) providing a first population of RNA molecules
20 each having a first region of random sequence;
- b) contacting the first population of RNA molecules with the first ligand;
- c) isolating a first ligand-binding subpopulation of the first population of RNA molecules by
25 partitioning RNA molecules in this first population which specifically bind the first ligand from those which do not;
- d) amplifying the first ligand-binding subpopulation in vitro;
- 30 e) identifying a first ligand binding sequence;
- f) preparing a second population of RNA molecules each of the RNA molecules including the first ligand binding sequence and a second region of random sequence;

- 12 -

g) contacting the second population of RNA molecules with a second ligand capable of binding the first ligand binding sequence; and

h) isolating a subpopulation of the catalytic
5 RNA molecules from the second population of RNA molecules by partitioning RNA molecules which have been modified in step g) from those which have not been modified.

In various preferred embodiments of the method, the first ligand is ATP, the first ligand is biotin, the
10 second ligand serves as a substrate for the chemical reaction, and the first and second ligands are the same.

In other preferred embodiments of the method, the catalytic RNA molecule can transfer a phosphate from a nucleotide triphosphate to the catalytic RNA molecule.
15 In more preferred embodiments of the method, the transfer is to the 5'-hydroxyl of the catalytic RNA molecule and the transfer is to an internal 2'-hydroxyl of the catalytic RNA molecule.

In another preferred embodiment of the method, the
20 catalytic RNA molecule can transfer a phosphate from a nucleotide triphosphate to a nucleic acid (preferably, a ribonucleic acid) other than the catalytic RNA molecule.

In another preferred embodiment of the method, the catalytic RNA molecules can catalyze N-alkylation, the
25 catalytic RNA molecule can catalyze N-alkylation of the catalytic RNA molecule, and the catalytic RNA molecule can catalyze N-alkylation of a nucleic acid other than the catalytic RNA molecule.

In another aspect, the invention features a
30 catalytic RNA molecule which can transfer a phosphate from a nucleotide triphosphate to the catalytic RNA molecule. In preferred embodiments, the transfer is to the 5'-hydroxyl of the catalytic RNA molecule and the transfer is to an internal 2'-hydroxyl of the catalytic
35 RNA molecule.

- 13 -

In another aspect, the invention features a catalytic RNA molecule which can transfer a phosphate from a nucleotide triphosphate to a nucleic acid (preferably, a ribonucleic acid) other than the catalytic
5 RNA molecule.

In another aspect, the invention features a catalytic RNA capable of catalyzing N-alkylation. In preferred embodiments, the catalytic RNA molecule can catalyze N-alkylation of the catalytic RNA molecule, and
10 the catalytic RNA molecule can catalyze N-alkylation of a nucleic acid other than the catalytic RNA molecule.

In another aspect, the invention features a method for producing a catalytic RNA molecule capable of binding a first ligand and catalyzing a chemical reaction
15 modifying a first substrate molecule bound to the catalytic RNA molecule. The method entails the following steps:

- a) providing a first population of RNA molecules each having a first region of random sequence;
- 20 b) contacting the first population with the first ligand;
- c) isolating a first ligand-binding subpopulation of the first population of RNA molecules by partitioning RNA molecules in the first population of RNA
25 molecules which specifically bind the first ligand from those which do not;
- d) amplifying the first ligand binding subpopulation in vitro;
- e) identifying a first ligand binding sequence;
- 30 f) preparing a second population of RNA molecules each of the RNA molecules including the first ligand binding sequence and a second region of random sequence, each of the RNA molecules being bound to the first substrate molecule;

- 14 -

g) contacting the second population of RNA molecules with a second ligand capable of binding the first ligand-binding sequence; and

h) isolating a subpopulation of the catalytic RNA molecules from the second population of RNA molecules by partitioning RNA molecules which are bound to a substrate molecule which has been modified in step g) from those RNA molecules which are bound to a substrate molecule which has not been modified in step g).

10 In a preferred embodiment of this method, the second ligand serves as a second substrate for the chemical reaction.

The invention also features ribozymes having polynucleotide kinase activity. Such ribozymes have 80%,
15 preferably 85%, more preferably 95% homology to any of classes I - VII polynucleotide kinase ribozymes described in FIG. 5. More preferably such ribozymes have 90% (more preferably 95%) homology to the core catalytic region of any of these classes of ribozymes. The core catalytic
20 region is the minimal sequence required for catalytic activity. This sequence can be determined using standard deletion analysis.

The invention also features ribozymes capable of carrying out an alkylation reaction. In a preferred
25 embodiment the ribozyme has 90%, and preferably 95% homology to BL-E.

Other features and advantages of the invention will be apparent from the description of the preferred embodiments, and from the claims.

30 Description of the Drawings

FIG. 1 is a schematic illustration of a minimal ATP aptamer (SEQ ID NO: 4).

FIG. 2 is a schematic illustration of the random RNA pool built around the ATP aptamer structure and the
35 selection scheme (SEQ ID NO: 5). The pool contained

- 15 -

three regions of random sequence (N) for a total of 100 randomized bases. The aptamer region was mutagenized to a level of 15%. The *Ban* I site used to ligate the two halves of the pool is shown in gray. Constant primer
5 binding sites are shown as thick lines. Random pool RNA was allowed to react with ATP- γ -S and thiophosphorylated molecules were isolated by reaction with thiopyridine-activated thiopropyl sepharose. Non-specifically bound
10 molecules were removed by washing under denaturing conditions. Active molecules were eluted with 2-mercaptoethanol. Constant regions: 5'-GGAACCUCUAGGUCAUUAAGA-3' (5'-end constant region) (SEQ ID NO: 1); 5'-ACGUCAGAAGGAUCCAAG-3' (3'-end constant region) (SEQ ID NO: 2).

15 FIG. 3 is a graph showing the percent RNA eluted by 2-mercaptoethanol from the thiopyridine-activated thiopropyl Sepharose at each cycle of selection. Background sticking and elution from the resin is approximately 0.5%. The concentration of ATP- γ -S used in
20 each selection and the incubation time for each selection is shown below the graph. Also indicated is whether the selection entailed mutagenic PCR.

FIG. 4 is a graph showing the k_{obs} of pool RNA for selection cycles 6-10, 12 and 13. Reactions were
25 performed with 100 μ M ATP- γ -S, and a time point was chosen such that < 20% of the pool had reacted. At cycle 6, the activity of the pool could be readily detected. The following seven cycles increased the activity by nearly three orders of magnitude. The drop in k_{obs} in
30 cycle 8 is presumably due to the effects of mutagenic PCR, coupled with the fact that the pool was no longer immobilized on streptavidin in this cycle. Cycle 11 activity declined for unknown reasons.

FIG. 5 illustrates the sequences of molecules
35 representing the seven major kinase classes (50 clones

- 16 -

sequenced) (SEQ ID NOS: 6-24). Arrows delimit the ATP aptamer conserved loop. The *Ban* I site used for pool construction (see FIG. 2) is underlined. Complementarity between the random region and the (constant) 5'-end of the RNA is shaded (Classes I and V). Both of these classes are 5'-kinases; these regions may serve to bind the 5'-end in the active site of the ribozymes. Sites of 2'-thiophosphorylation are shown as white letters in black boxes. Clone Kin. 47 is inactive, and contains a G to A mutation at the site of 2'-thiophosphorylation. The sequences of the constant primer binding regions (see FIG. 2) are not shown except for the first three bases following the 5' primer binding site (AGA). The length of the original pool (not including primer binding sites) was 138 nucleotides. Point deletions may have occurred during the chemical synthesis of the pool DNA, and larger deletions may be due to annealing of primers to sites in the random regions during reverse-transcription or PCR.

FIG. 6 is a set of schematic illustrations of proposed structures of the ATP aptamer consensus and several classes of ATP aptamer (SEQ ID NOS: 25-29). In the illustration of the consensus aptamer conserved bases in the loop are shown in capital letters. Positions that tend to be A, but which can vary, are shown as "a"s. The bulged G is also conserved, but the stem regions (aside from being base paired) and the right hand loop are not. For the schematic illustrations of possible secondary structures of the ATP aptamer domains of four of the major classes of ribozymes, the sequence of the most active clone is shown in each case. Positions in the loop regions that differ from the consensus sequence for the ATP aptamer are highlighted in gray. One of the stem regions from each of Classes II, VI and VII is missing, and so these structures are not shown.

- 17 -

FIG. 7A is a schematic illustration of a ribozyme capable of transferring a phosphate to its 5' end (SEQ ID NO: 30). FIG. 7B is a schematic of a trans-acting ribozyme and a substrate (GGAACCU).

5 FIG. 8A is a strategy for *in vitro* evolution of self-alkylating ribozymes. FIG. 8B is a scheme for isolating biotin-binding RNAs by affinity chromatography. FIG. 8C is a scheme for isolating self-biotinylation RNA enzymes. FIG. 8D shows coding sequences for RNA pools
10 used for *in vitro* selection experiments (SEQ ID NOS: 32-34). Upper case A, C, G, T: pure nucleotide. N: equimolar mix of A, C, G, T. Lower case a, c, g, t: 70% major nucleotide, 10% each of three minor nucleotides. Underline: constant primer sequences used for
15 amplification.

FIG. 9A illustrates progress of the biotin aptamer selection. Biotin-eluted RNA expressed as a percentage of total RNA applied to the biotin-agarose column is plotted as a function of selection cycle. Individual
20 RNAs eluted from the seventh round were subcloned and sequenced. Greater than 90% of the clones correspond to the sequence shown in FIG. 12. FIG. 9B illustrates progress of the self-biotinylation selection. Ligation rate determined by incubation with 200 μ M BIE followed by
25 streptavidin-agarose purification. Values are corrected for 0.02% non-specific RNA binding.

FIG. 10A is a site-specific alkylation reaction catalyzed by BL8-6 ribozyme. 5'-end labeled BL8-6 RNA was allowed to react overnight with 200 μ M BIE and then
30 separated by streptavidin affinity chromatography into biotinylated and non-biotinylated fractions. RNA was then treated with sodium borohydride and aniline acetate to specifically cleave at N7 alkylation sites. For comparison, DMS-treated RNA was treated in parallel. A
35 single major cleavage site in the biotinylated fraction

- 18 -

(corresponding to Gua-96) is absent from the non-biotinylated RNA. Minor bands present in the non-biotinylated fraction appear to result from non-specific RNA cleavage as judged by their greater intensity in BL8-5 6 RNA subjected to partial alkaline hydrolysis. FIG. 10B illustrates the inferred N-alkylation reaction at the N7 position of G-96.

FIG. 11 illustrates functional biotin binder and biotin ligator sequences (SEQ ID NOS: 35-86). The 10 partially-randomized pool sequence is shown above each set of sequences. Deviations from the principle nucleotide at each position are explicitly written while conservation of the wide type base is indicated with a dash. Biotin aptamer and self-biotinylating RNA 15 partially-randomized pools were re-selected for biotin-agarose binding and self-biotinylation respectively. Biotin aptamer sequences correspond to clones from the fourth round of re-selection. Self-biotinylating ribozyme clones were sequenced after eight rounds of re- 20 selection, when the overall biotinylation activity of the pool was 100 times the activity of the initial BL8-6 ribozyme. Arrows are used to indicate the locations of proposed helices. Boxed nucleotides are highly conserved yet not involved in secondary structure.

25 FIG. 12A and FIG. 12B illustrate the proposed secondary structures for the biotin aptamer and the self-biotinylating ribozyme. Nucleotides within the boxed region are highly conserved and make up the essential core of the aptamer and ribozyme. Asterisks indicate 30 pairs of positions that co-vary in a Watson-Crick sense. Nucleotides in the constant primer sequences are shown in italics. FIG. 12A is a complete sequence of the BB8-5 biotin aptamer, shown as the proposed pseudoknot. FIG. 12B (SEQ ID NO: 91) is a sequence and proposed cloverleaf 35 structure for the BL8-6 self-biotinylating ribozyme. The

- 19 -

guanosine residue that serves as the alkylation site for the biotinylation reaction is circled.

FIG. 13A illustrates the sequence (SEQ ID NO: 87) of a clone obtained from the partially-randomized
5 ribozyme pool after re-selecting for biotinylation activity (BL2.8-9) was modified to allow folding into an idealized cloverleaf structure. FIG. 13B illustrates *in vitro* transcribed RNA assayed for self-biotinylation with 10 μ M BIE. Folding was calculated by the LRNA Program
10 (Zuker, Science 244:48, 1989)

FIG. 14B shows the results of a ribozyme-mediated biotinylation of a separate RNA substrate. The designed self-biotinyllating ribozyme (FIG. 13A) was re-engineered into two halves, BL-E and BL-S, that could respectively
15 serve as the enzyme and substrate for the biotinylation reaction. This re-engineered molecule is illustrated in FIG 14A (SEQ ID NOS: 88, 89). To assay the two piece system, 5 μ M radiolabelled BL-S RNA was incubated in the presence of 200 μ M BIE and 0 to 500 nM unlabelled BL-E
20 RNA. RNA biotinylation was determined as described herein. The reaction plateaus overnight at a level corresponding to one equivalent of product.

Description of the Preferred Embodiments

EXAMPLE 1

25 In one example of the invention, RNA molecules which bind ATP were first isolated from a pool of random RNA. RNA molecules capable of binding ATP were sequenced, and the information obtained was used to design a second pool of RNA molecules which included an
30 ATP binding site or variant thereof. This pool was then subjected to selection and amplification to identify RNA molecules having polynucleotide kinase activity.

- 20 -

Selection of ATP-binding RNAs: The selection of ATP-binding RNAs was carried out in a manner designed to ensure selection of RNAs capable of binding ATP in solution as well as on an insoluble support. This was
5 accomplished by selecting RNA molecules which bound an ATP-sepharose column and could be eluted using ATP.

A chemically synthesized pool of DNA molecules containing a central region of 120 random nucleotides flanked by constant regions used as primer binding sites
10 was PCR-amplified and transcribed *in vitro* by T7 RNA polymerase in the presence of [α -³²P]GRP. RNA was ethanol-precipitated and unincorporated nucleotides removed by Sephadex-G50 chromatography. Following a brief incubation at 65°C in binding buffer (300 mM NaCl,
15 20 mM Tris, pH 7.6, 5 mM MgCl₂), the RNA was cooled to room temperature before being loaded onto a 1-ml ATP agarose affinity column. The affinity matrix contained ~1.6 mM ATP linked through its C8 position through a diaminohexyl linker to cyanogen bromide-activated agarose
20 (Sigma, St. Louis, MO). After washing with 6 column-volumes of binding buffer, bound RNAs were eluted with 3 column-volumes of binding buffer containing 4 mM ATP, then concentrated by precipitation with ethanol. For the first three cycles, an agarose pre-column was used to
25 prevent enrichment of the RNA pool with agarose-binding RNAs, and bound RNAs were eluted with 5 mM EDTA in water rather than affinity-eluted with ATP. After reverse transcription and PCR amplification, DNA templates were transcribed and the resulting RNA was used in the next
30 round of selection. RNA from the eighth round of selection was converted to cDNA, amplified as double-stranded DNA by PCR, digested with *Eco*R1 and *Bam*H1, gel-purified and cloned into the phage M13 based vector pGem3Z (Promega, Madison, WI).

- 21 -

Thirty-nine clones from the eighth cycle RNA population were sequenced seventeen different sequences were found. Of these, the most abundant sequence (C8-ATP-3) occurred 14 times, and 12 sequences occurred just once. Comparison of the seventeen different sequences revealed an 11-nucleotide consensus sequence, of which seven positions are invariant among all clones but one (C8-ATP-15). Clones 2, 3, 8, 15, and 19 were individually tested for binding to ATP-agarose. All had a dissociation constant (K_d) of less than 50 μ M, except for C8-ATP-15, for which the estimated K_d was ~250 μ M.

To determine the minimal sequence for ATP binding, deletions of C8-ATP-3 were analyzed. An active RNA molecule 54 nucleotides in length (ATP-54-1) was generated by a combination of 5' and 3' deletions. This RNA can be folded into a secondary structure in which the 11-base consensus is flanked by two base-paired stems. Deletion of the left-hand stem abolished ATP-binding activity; dimethylsulphate modification experiments also supported the proposed secondary structure. Comparing sequences of all the clones showed that they all had a potential to fold into this secondary structure. This analysis also highlighted the presence of an invariant unpaired G opposite the 11-base consensus. The orientation and distance of this G and its flanking sequences relative to the consensus sequence was variable from clone to clone. The stems flanking the conserved G and the consensus were variable in length and composition, and frequently contained G-U base pairs. The simplest explanation for the observation that all of the selected clones contained a single consensus sequence embedded in a common secondary structure is that these clones contain the shortest sequences capable of binding ATP with the necessary affinity, and that all other sequences with comparable or superior affinity are longer

- 22 -

and hence less abundant in the initial random sequence pool.

On the basis of these findings, a smaller RNA of 40 nucleotides (ATP-40-1) was designed, in which the consensus sequence was flanked by stems of six base pairs, with the right-hand stem closed by a stable loop sequence for enhanced stability. This RNA bound ATP as well as the full-length 164-nucleotide RNA C8-ATP-3 and was used for later experiments. Variant 40-oligonucleotide RNAs were also synthesized to test the importance of the highly conserved unpaired G (residue G34 in ATP-40-1) for ATP binding. Deleting this residue or changing it to an A residue eliminated binding, confirming the results of the selection experiments.

To determine which functional groups on the ATP are recognized by the ATP-binding RNA, we examined the ability of a series of ATP analogues to elute bound ATP-40-A RNA from an ATP-agarose column. Methylation of positions 1, 2, 3, or 6 on the adenine base, or the 3' hydroxyl of the ribose sugar, abolish binding, as does removal of the 6-amino or 2'-hydroxyl. Positions 7 and 8 on the base can be modified without effect; this is not surprising considering that selection was for binding to ATP linked to an agarose matrix through its C8 position. Adenosine, AMP, and ATP are equally efficient at eluting the RNA, suggesting that the 5' position on the ribose moiety is not recognized by the RNA.

Isocratic elution (Arnold et al., J. Chromatography 31:1, 1986) from ATP-agarose and equilibrium gel filtration (Fersht, in Enzyme Structure and Mechanism p. 186-188, Freeman, New York, 1985) was used to measure the dissociation constant for the RNA-ATP complex on the column and in solution. The K_d of ATP-40-1 was $\sim 14 \mu\text{M}$ when measured by isocratic elution from an ATP-C8-agarose column, and $6-8 \mu\text{M}$ by equilibrium gel

- 23 -

filtration. The K_d for the ATP-agarose complex is an upper estimate, because the fraction of bound ATP that is accessible to the RNA is not known. The solution K_d for adenosine was similar to that of ATP (5-8 μM), but the K_d for dATP was not measurable ($>1 \text{ mM}$). The K_d of ATP-40-1 for its ligand dropped to 2 μM when the Mg^{+2} concentration was raised from 5 to 20 mM. Changing the base pair U18-A33 to C-G, as found in most of the clones initially selected, further decreased the K_d to 0.7 μM . At almost saturating concentrations of ATP (50 μM), the RNA bound ~0.7 equivalents of ATP. The RNA likely binds its ligands with a stoichiometry of unit.

Kethoxal modification (Moozod et al., J. Mol. Biol. 187:399, 1987) was used to assess the accessibility of guanosine residues to modification. G7 and G17 within the loop, and G6 (which forms the G-C base pair on the left side of the loop), all of which are strongly protected in the absence of ATP, become highly accessible to modification by this reagent in the presence of ATP. Other guanosine residues, including G8 in the large loop, the single unpaired G opposite the loop, and Gs in the stems, are highly protected in the presence or absence of ATP. These observations suggest that the motif is highly structured both in the presence and absence of ATP, but that binding induces a conformational change in the structure of the RNA.

A pool of random sequence RNAs, using the above-identified minimal ATP aptamer as a core structure was prepared and used to create polynucleotide kinase ribozymes. The ATP aptamer is based on that described by Sassanfar and Szostak (Nature, 364:550, 1993).

Selection of Catalytic RNAs: A pool of RNA molecules for selection of catalytic RNAs was created based on a minimal ATP aptamer core sequence (FIG. 1). The ATP aptamer core was surrounded by three regions of random

- 24 -

sequence, 40, 30, and 30 nucleotides in length as shown in FIG. 2. The ATP-binding domain itself was mutagenized such that each base had a 15% chance of being non-wild type, to allow for changes in the aptamer sequence that might be required for optimal activity. To increase the likelihood of finding active molecules, an effort was made to create a pool containing as many different molecules as possible. Because it is difficult to obtain an acceptable yield from the synthesis of a single oligonucleotide of this length (174 nucleotides), two smaller DNA templates were prepared and linked together to generate the full length DNA pool (FIG. 2) (Bartel and Szostak, Science, 261:1411, 1993). The presence of constant primer binding sites at the 5' and 3' ends of the molecules permitted amplification by PCR. Transcription of this DNA pool yielded between 5×10^{15} and 2×10^{16} different RNA molecules.

In order to select for catalytic activity, it is necessary to tag active molecules so that they can be separated from inactive ones. To accomplish this, the random sequence RNA pool was incubated with ATP- γ -S and the transfer of the thiophosphate from ATP- γ -S to the RNA was selected for chromatography on a thiopyridine-activated thiopropyl sepharose column, which forms disulfide bonds with RNAs containing thiophosphate groups. Molecules without thiophosphates were washed away under denaturing conditions. RNAs linked via a disulfide to the column matrix were eluted with an excess of 2-mercaptoethanol. This overall scheme is illustrated in FIG. 2. Briefly, the pool was incubated with ATP- γ -S under conditions designed to promote the formation of RNA tertiary structure (400 mM KCl, 50 mM MgCl₂, 5mM MnCl₂, 25mM HEPES, pH 7.4). Mn²⁺ was included because of its ability to coordinate phosphorothioates. Streptavidin agarose immobilization of pool RNA was used during the

- 25 -

first seven cycles to prevent pool aggregation. After cycle 7, the ATP- γ -S reaction step was performed in solution (1 μ M RNA). For the first cycle, 2.4 mg (40 nmoles; 5 pool equivalents) of random pool RNA was used, in the second cycle 150 μ g (2.4 nmoles) RNA was used, and in succeeding cycles 60 μ g (1 nmole) was used. The selection step was performed by incubating the RNA with thiopyridine-activated thiopropyl sepharose-6B (Pharmacia, Piscataway, NJ) in 1 mM EDTA, 25 mM HEPES, pH 7.4 for 30 minutes at room temperature. The resin was then washed with 20 column volumes each of wash buffer (1M NaCl, 5 mM EDTA, 25 mM HEPES, pH 7.4), water, and finally 3 M urea, 5 mM EDTA to eliminate molecules without thiophosphates. RNAs linked to the resin via a disulfide were eluted with 0.1 M 2-mercaptoethanol in 0.5X wash buffer. Reverse-transcription, PCR and transcription yielded a new RNA pool enriched in active molecules. This process comprised one cycle of selection.

Prior to each cycle of the selection, the pool RNA generated by transcription was exhaustively dephosphorylated with calf intestinal alkaline phosphatase to remove the 5'-triphosphate, and any other phosphates that might have been transferred to the RNA by autophosphorylation during transcription.

The selection protocol demanded only that an RNA molecule contain a thiophosphate in order for it to be isolated. Reactions that could have been selected for include: transfer of the γ -thiophosphate from ATP- γ -S to the 5'-hydroxyl of the RNA (analogous to the reaction catalyzed by T4 polynucleotide kinase), to the 3'-end of the RNA, to an internal 2'-hydroxyl, or even to a group on one of the bases. Transfer of diphosphate (or perhaps the entire triphosphate) instead of a single thiophosphate is also possible for all of these

- 26 -

reactions. A splicing reaction, in which ATP- γ -S displaces one of the first few nucleotides of the RNA in a manner analogous to the reaction catalyzed by the Group I introns, could also occur. However, cleavage of more than the first few bases of the RNA would result in a molecule lacking a 5'-primer binding site, and such a molecule would not be amplified during the PCR step of the selection. Similarly, any reaction that blocked reverse transcription would not be selected for.

10 The progress of the selection process was monitored by measuring the fraction of the pool RNA that bound to the thiopropyl Sepharose and was eluted with 2-mercaptoethanol (FIG. 3). Initially, ~0.5% of the RNA bound nonspecifically to the matrix and was eluted by 2-mercaptoethanol. After five cycles of selection, greater than 20% of the pool RNA reacted with thiopropyl Sepharose. Since there were at least 10,000 different molecules left in the pool at this stage, the stringency of the selection in the succeeding cycles was increased by lowering the ATP- γ -S concentration and the incubation time, in order to try to isolate the most active catalysts.

Optimization of Catalytic RNAs: Because the random pool initially prepared sampled sequence space very sparsely (there are between 4^{100} and 10^{60} possible 100-mers, but only approximately 10^{16} different molecules in the pool), active molecules are likely to be sub-optimal catalysts. Accordingly, three cycles of mutagenic PCR (before selection cycles 7, 8, and 9) were performed to allow the evolution of improvements in the active molecules. Mutagenic PCR was performed as described by Bartel and Szostak (Science, 261:1411, 1993) and by Cadwell and Joyce (PCR Methods Appl., 2:28, 1992). Briefly, thirty total cycles of PCR were done at each round to yield ~ 2% mutagenesis. Reactions of pool RNAs were performed

- 27 -

either with trace ATP- γ -³⁵S, or with 100 μ M ATP- γ -S plus additional trace ATP- γ -³⁵S. Dithiothreitol (DTT, 10 mM) was included in the reactions. Reactions were quenched by the addition of one volume of 150 mM EDTA, 20 mM DTT in 95% formamide. Reactions were analyzed by electrophoresis on 10% polyacrylamide/8 M urea gels. Quantitation was performed using a PhosphorImager (Molecular Dynamics). A known amount of ATP- γ -³⁵S was spotted on the gels as a standard. The combined effect of increasing the stringency and performing mutagenic PCR was to increase the activity of the pool by nearly three orders of magnitude from cycle 6 to cycle 13 (FIG. 4). Catalytic RNAs Identified: After 13 cycles of selection, RNA molecules from the pool were cloned using the pT7 Blue T-Vector kit by Novagen, and 50 clones were sequenced. The clones sequenced (FIG. 5) fall into seven classes of two or more closely related molecules (19 clones) and 31 unique sequences. Each class of sequences represents molecules with a common ancestor that acquired mutations during the course of the mutagenic PCR done in cycles 7-9 of the selection.

Comparison of the sequences in the seven major classes of molecules reveals significant conservation of the sequence of the original ATP binding site in some of the active RNAs. FIG. 6 shows the putative structures for the ATP aptamer regions from Classes I, III, IV and V, the classes for which an aptamer-like structure can be drawn. It appears that Classes I and III have changed significantly from the original ATP binding domain, whereas Classes IV and V are only slightly different from the ATP aptamer consensus sequence described by Sassanfar and Szostak (Nature, 364:550, 1993). Either the right or left hand stems of the Class II, VI and VII aptamer regions appear to be missing, and it seems likely that these molecules have found novel modes of binding their

- 28 -

substrates. Using run-off transcription of synthetic DNA oligonucleotides (Milligan and Uhlenbeck, Methods Enzymol. 180:51, 1989) the RNAs corresponding to the Class I, III, IV, and V aptamer regions were produced.

- 5 The Class IV aptamer RNA binds weakly to C-8 linked ATP agarose (Sassanfar and Szostak, supra), consistent with a molecule having a K_d for ATP in the range of 0.05-0.5 mM. The Class I, III, and IV aptamers, on the other hand, do not detectably interact with ATP agarose, consistent with
- 10 K_d s > 0.5 mM for ATP (if they bind ATP at all).

Presumably, the corresponding classes of kinases have developed novel modes of binding ATP- γ -S.

- Characterization of the Catalyzed Reactions: Pool 13 RNA and the members of each of the major classes of kinases
- 15 were tested to determine what reactions they catalyze. Nuclease P1 analysis was performed as follows. RNA (1 μ M) was allowed to react with ~1 μ M ATP- γ -³⁵S in reaction buffer for 4-18 hours. The RNA was separated from nucleotides by G-50 spin column gel filtration
- 20 (Boehringer-Mannheim, Indianapolis, IN). The RNA was digested with nuclease P1 as described in Westaway et al. (J. Biol. Chem. 268:2435, 1993) and Konarska et al. (Nature 293:112, 1981). An aliquot was then spotted directly onto a PEI cellulose TLC plate (Baker,
- 25 Phillipsburg, NJ) and developed in 1M LiCl, 10mM DTT (as described in Westaway, supra). The products were localized by UV shadowing (for unlabelled GMP α S) or autoradiography. Thiophosphate containing nucleotides run slower in this system than do the corresponding
- 30 phospho-nucleotides, presumably because there is weaker interaction between Li⁺ and the thiophosphate than there is with the phosphate.

PEI cellulose thin layer chromatography (TLC) of nuclease P1 digests of auto-thiophosphorylated RNA shows

35 two major radiolabeled products, demonstrating that at

- 29 -

least two different reactions are catalyzed by the pool
13 RNAs. If a particular RNA molecule transfers the γ -
thiophosphate from ATP- γ -S to its own 5'-hydroxyl, the
nuclease P1 digestion should yield labeled GMP α S, since
5 all of the RNAs begin with guanosine. All members of
Classes I, II, III, V, and VI yield GMP α S as the sole
nuclease P1 digestion product, indicating that they are
5'-kinases. Classes IV and VII, on the other hand, yield
a nuclease P1 digestion product that does not migrate
10 from the origin in the TLC system used. Both RNase T2,
which hydrolyzes RNA to nucleotide 3'-monophosphates, and
nuclease P1 digestion of reacted Class IV and Class VII
RNAs, give products that run as molecules with charges of
-5 to -6 on DEAE cellulose TLC plates, using a solvent
15 system that separates based upon the charge of the RNA
fragment (Dondey and Gross, Anal. Biochem. 98:346, 1979;
Konarska et al., Nature 293:112, 1981). These data are
consistent with Class IV and VII RNAs being internal 2'-
kinases, since neither nuclease P1 nor RNase T2 can
20 cleave at 2'-phosphorylated sites (Westaway et al.,
J. Biol. Chem. 268:2435, 1993). The products of these
digestions, then, should be 35 S-labeled dinucleotides with
5'-phosphates or 3'-phosphates (for nuclease P1 and RNase
T2 digestions, respectively) and 2'-mono- or di-
25 phosphates.

Experiments in which the RNAs were allowed to
react with unlabeled ATP- γ -S and were then purified and
reacted with ATP- γ - 32 P and T4 polynucleotide kinase
support the proposal that Classes I, II, III, V, and VI
30 are 5'-kinases, and that Classes IV and VII phosphorylate
some internal site. As expected, reaction products from
Classes I, II, III, V, and VI cannot be labeled by T4
polynucleotide kinase, consistent with their being 5'-
kinases. Class IV and VII RNAs, on the other hand, are
35 efficiently labeled by T4 polynucleotide kinase after

- 30 -

they have been allowed to react with ATP- γ -S.

Furthermore, this labeled RNA can be purified on a thiopyridine-activated thiopropyl sepharose column, demonstrating that the thiophosphate label is not lost
5 during the reaction with ATP and T4 polynucleotide kinase. Thus, the Class IV and VII kinases do not catalyze reactions involving their 5'-hydroxyls.

Conclusive evidence for the 2'-kinase hypothesis is provided by partial alkaline hydrolysis of the auto-
10 thiophosphorylated, 5'-³²P-labeled RNA. For this analysis, RNA was reacted with ATP- γ -S as described above for TLC analysis, except that 100 μ M unlabeled ATP- γ -S was used. The thiophosphorylated RNAs were purified on thiopyridine-activated thiopropyl sepharose, and then 5'-
15 end labeled using T4 polynucleotide kinase and ATP- γ -³²P. Alkaline hydrolysis was performed in 50 mM sodium carbonate/bicarbonate buffer, pH 9.0, 0.1 mM EDTA for 3 min. at 90°C. Reaction products were analyzed on an 8% polyacrylamide/8 M urea gel.

20 For RNAs from both Classes IV and VII, a gap is seen in the alkaline hydrolysis ladder of the auto-thiophosphorylated material that is not present in the ladder made with unreacted RNA. The missing bands can be most easily explained if the 2'-hydroxyls at these
25 positions are thiophosphorylated, thus preventing base-catalyzed RNA hydrolysis. This experiment permitted identification of positions of thiophosphorylation: G62 in Kin.10 (Class IV) and G83 in Kin.62 (Class VII). G62 is in a putative helix within the ATP aptamer region of
30 Kin.10, and G83 is in the random loop between the two halves of Kin.62's aptamer domain.

Kinetic Analysis of Kinase Ribozymes: Kinetic analysis of the most active clone from each of the four major classes of kinases has revealed that they all obey the
35 standard Michaelis-Menten kinetics expected of molecules

- 31 -

possessing saturable substrate binding sites. Rates for each clone were determined (as described herein) at 6 different ATP- α -S concentrations, ranging from 2 μ M - 2.5 μ M. Values of k_{cat} and K_m are shown in Table 1, and range 5 between 0.03 and 0.37 min^{-1} and between 41 and 456 μ M, respectively.

TABLE 1

	<u>Kinase Class (Clone)</u>	<u>k_{cat} (min^{-1})</u>	<u>K_m (μM)</u>
10	Class I (Kin.46)	0.37 \pm 0.01	456 \pm 57
		0.23 \pm 0.02	116 \pm 41
		0.36 \pm 0.02	352 \pm 85
15	Class II (Kin.25)	0.20 \pm 0.02	41 \pm 15
		0.33 \pm 0.02	42 \pm 11
	Class III (Kin.42)	0.07 \pm 0.005	50 \pm 13
		0.10 \pm 0.016	58 \pm 28
	Class IV (Kin.44)	0.03 \pm 0.001	276 \pm 25
		0.03 \pm 0.001	200 \pm 22

The k_{cat} for Class I-IV ribozymes compares favorably with corresponding values for naturally occurring ribozymes, 20 which range from 0.04 to 2 min^{-1} . Comparison of k_{cat}/K_m is difficult because most natural ribozymes have oligonucleotide substrates that form base pairs with the ribozyme's substrate binding site, leading to very low K_m values. A comparison between the kinase ribozymes 25 described here and the self-cleavage reaction catalyzed by the *Tetrahymena* Group I intron is particularly relevant, however, because both reactions use external small molecule substrates (ATP- γ -S and guanosine nucleotides, respectively) to modify themselves. Kin.25 30 (Class II) phosphorylates itself with a k_{cat} of approximately 0.3 min^{-1} and a k_{cat}/K_m of $6 \times 10^3 \text{ min}^{-1} \text{ M}^{-1}$. The *Tetrahymena* self-splicing intron has a k_{cat} of 0.5 min^{-1} and a k_{cat}/K_m of $2.5 \times 10^4 \text{ min}^{-1} \text{ M}^{-1}$ (Bass and Cech,

- 32 -

Nature 308:820, 1984). Thus, from a vanishingly small sampling of sequence space, it has been possible to isolate a molecule with autocatalytic activity essentially as good as that of a ribozyme found in nature.

Class I-IV kinases show specificity for ATP- γ -S as a substrate. No reaction (<0.1% ATP- γ -S rate) could be detected with GTP γ S, indicating that the RNAs can discriminate between similar substrates. Interestingly, as much as 30% of the cycle 13 pool RNA can use GTP- γ -S as a substrate, and thus pool 13 does contain molecules with less stringent substrate specificities. The Class I-IV kinases are also able to discriminate between ATP- γ -S and ATP ($k_{\text{obs}}(\text{ATP-}\gamma\text{-S})/k_{\text{obs}}(\text{ATP})$: Class I = 55; Class II = 300; Class III = 150; Class IV \geq 300; 100 μ M ATP, ATP- γ -S). Since these values are significantly larger than the three to ten fold intrinsic reactivity difference between ATP- γ -S and ATP (Herschlag et al., Biochemistry 30:4844, 1991), the data suggest that the thiophosphate is important for binding, catalysis or both. Furthermore, pool 13 RNA is not detectably labeled by either ATP- α - ^{35}S or ATP- α - ^{32}P , suggesting that 5' splicing is not a reaction that occurs in the pool (unless the γ -thiophosphate is an absolute requirement for the molecules that carry out this reaction).

Rate Acceleration: The uncatalyzed background reaction for the thiophosphorylation of RNA (or guanosine) by ATP- γ -S was not detectable. Based on the sensitivity of these experiments, the lower limit for the rate acceleration of the kinase ribozymes is roughly 10^5 -fold. At 70°C the rate of hydrolysis of ATP in the presence of Mg^{2+} is $\sim 4 \times 10^{-4} \text{ min}^{-1}$ (pH 6-8). Correcting for the temperature and 55 M water, this value gives a second order rate constant of approximately $1 \times 10^{-6} \text{ min}^{-1} \text{ M}^{-1}$. ATP- γ -S should hydrolyze 3-10 times faster than ATP.

- 33 -

Taking this factor into account, the approximate rate enhancement of the present ribozymes

$[k_{\text{cat}}/K_m]/[k_{\text{hydrolysis}}]$, would be $6 \times 10^3 \text{ min}^{-1} \text{ M}^{-1}/\sim 10^{-5} \text{ min}^{-1} \text{ M}^{-1}$ or $10^8 - 10^9$ fold. This enhancement corresponds

5 to an effective molarity of $10^4 - 10^5 \text{ M}$ for ATP in the ATP-ribozyme complex ($k_{\text{cat}}/k_{\text{hydrolysis}} = 0.3 \text{ min}^{-1}/10^{-5} \text{ min}^{-1} \text{ M}^{-1}$). A comparison of first-order rate constants gives a value for the rate enhancement that is independent of substrate binding. This value is approximately 10^3 fold
10 ($k_{\text{cat}}/k_{\text{hydrolysis}}$ (1° order) = $0.3 \text{ min}^{-1}/\sim 4 \times 10^{-4} \text{ min}^{-1}$). This analysis assumes that the mechanism of hydrolysis of ATP- γ -S (dissociative) is the same as that used by the kinase ribozymes.

Intermolecular Catalysis and Turnover: At least one of
15 the selected kinases is capable of catalyzing the phosphorylation of a separate RNA substrate. In particular, Kin.46 (Class I) was demonstrated to transfer the γ -thiophosphate from ATP- γ -S to the 5'-end of a 6-mer oligoribonucleotide with the same sequence as the 5'-end
20 of the ribozyme. To carry out this experiment, RNA was incubated as described in FIG. 2 except that 2.5 mM ATP- γ -S was used, and 100 μM 5'-HO-GGAACC-3' RNA was added. The 6-mer was synthesized by run-off transcription (Milligan et al., Meth. Enzymol. 180:51, 1989) and was
25 dephosphorylated with calf intestinal alkaline phosphatase prior to ion-exchange HPLC purification. The thiophosphorylated 6-mer marker was made by end-labelling 5'-GGAACC-3' with ATP- γ - ^{35}S using T4 polynucleotide kinase. Products were analyzed on 20% acrylamide/8 M
30 urea gels. Full-length Kin.46 was found to catalyze the reaction approximately 500-fold more slowly than the autocatalytic reaction. Part of the reason for the decreased activity is likely to be competition for the active site between the 5'-end of the RNA and the
35 exogenous 6-mer substrate. When the 5'-constant region

- 34 -

of the RNA is removed (via PCR with an internal 5'-primer, followed by transcription), the activity increases ~100-fold, but is still 6 fold below that of the auto-thiophosphorylation reaction. (At saturating concentrations of 6-mer (100 μ M) and ATP- γ -S (2.5 mM) the initial rate of thiophosphorylation is 0.05 μ M/min with 1 μ M ribozyme. In comparison, the rate of auto-thiophosphorylation for full length Kin.46 RNA (1 μ M) with 2.5 mM ATP- γ -S is 0.3 μ M/min.) At 25°C the ribozyme performs approximately 60 turnovers in 24 hours, and is thus acting as a true enzyme. The cause of the lower trans activity relative to the autocatalytic activity remains unknown, but could involve slow substrate binding or improper folding of the shortened ribozyme. The off rate of the 6-mer is not limiting because no burst phase is observed in a time course of the reaction.

The identification of autocatalytic ribozymes capable of carrying out catalysis in trans, i.e., catalyzing a reaction involving the ligand and a molecule other than the ribozymes itself can be found by testing the ability of the ribozyme to act on a molecule having a sequence similar to the region of the ribozyme which is modified.

FIG. 7A illustrates an example of a cis-acting ribozyme with polynucleotide kinase activity. A ribozyme capable of carrying out this catalysis in trans can be made by eliminating the 5' end of the ribozyme which would otherwise base pair with the 3' end of the ribozyme and be kinased. The particular molecule shown in FIG. 7B is derived from the molecule illustrated in FIG. 7A and transfers phosphate to the 5' end of the short oligoribonucleotide GGAACCU.

- 35 -

EXAMPLE 2

In a second example of the invention, RNAs which bind biotin were first created, identified, and isolated using a randomized RNA pool. The selected RNAs were used to prepare a second pool of partially randomized RNAs. This pool was then subjected to selection and amplification to identify RNAs capable of ligating biotin. The overall scheme is illustrated in FIGS. 8A, 8B, and 8C.

Selection of biotin-binding RNAs: A pool of approximately 5×10^{14} different random sequence RNAs was generated by *in vitro* transcription of a DNA template containing a central 72-nucleotide random sequence region, flanked at both ends by 20-nucleotide constant regions. This pool (random N72 pool) had the following sequence: GGAACACTATCCGACTGGCA(N)₇₂CCTTGGTCATTAGGATCG (SEQ ID NO: 3) (FIG. 8D, also SEQ ID NO: 32). On average, any given 28 nucleotide sequence has a 50% probability of being represented in a pool of this complexity. The initial pool of RNA (approximately 80 μ g; on average, 2-3 copies of each sequence) was resuspended in a binding buffer containing 100 mM KCl, 5 mM MgCl₂, and 10 mM Na-HEPES, pH 7.4, conditions chosen to favor RNA folding and to mimic physiological environments while minimizing non-specific aggregation. The solution was applied to an agarose column derivatized with 2-6 mM biotin (Sigma, St. Louis, MO) and subsequently washed with 15 column volumes of binding buffer. Specifically-bound RNAs were then eluted by washing the column with binding buffer containing 5 mM biotin. Ten μ g of glycogen and 0.3 M NaCl were then added to the eluted material, and the RNA was amplified as follows. Briefly, the mixture was precipitated with 2.5 volumes of ethanol at -78°C. After resuspending the selected RNA, the reverse transcriptase primer (2.5 μ M)

- 36 -

was annealed at 65°C for 3 min., and reverse transcription (RT) was carried out at 42°C for 45 min. (using Superscript RT enzyme, Life Technologies, Inc.). PCR amplification was performed by diluting one-fifth of the RT reaction with the appropriate dNTPs, PCR buffer, USB Taq polymerase (United States Biochemical, Cleveland, OH), and 0.5 μ M (+) primer containing the T7 RNA polymerase promoter. A strong band of the correct size was typically observed after 8-15 cycles amplification (94°C, 1 minute; 55°C, 45 seconds; 72°C, 1 minute). Half of the PCR reaction was used for *in vitro* transcription with T7 RNA polymerase (37°C, overnight). The resulting RNA was purified by electrophoresis on an 8% polyacrylamide gel.

After six rounds of repeated enrichment, more than half of the RNA applied to the biotin column was retained during the buffer wash, but eluted during the biotin wash (FIG. 9A). The RNA pool from the eighth round of selection was cloned into the PCR vector using the TA cloning kit (In Vitro-Gen, Inc., San Diego, CA), and individual aptamers were sequenced by the Sanger dideoxynucleotide method using the universal M13 primer. A single sequence (represented by clone BB8-5) accounted for >90% of the selected pool (two minor clones account for the vast majority of remaining RNAs).

Previous RNA selections for binding to small ligands, including various dyes, amino acids, cofactors, and nucleotides, have suggested that aptamers exist at a frequency of 10^{-10} to 10^{-11} in random sequence pools. All of these ligands, however, have contained aromatic rings which could intercalate between RNA bases and/or charged groups which might interact electrostatically with the RNA backbone. The lower frequency of biotin bindings (10^{-15}) shows that ligands lacking such groups may require a more complex binding site.

- 37 -

Selection for biotin-utilizing ribozymes: The sequence of the biotin aptamer was used to direct the synthesis of a second pool of RNAs which was screened for the presence of biotin-utilizing ribozymes (FIG. 8A). This pool
5 contained a core of 93 nucleotides (71 nucleotides derived from the original random region plus its 22 nucleotide 5' constant region; FIG. 8D) with the wild-type nucleotide (i.e., that which was found in the original biotin aptamer incorporated at each position in
10 the template with 70% probability (the three non-native nucleotides each occurring with 10% probability). Deletion analysis indicated that the 3' primer was not required for binding and the same sequence was therefore used for the 3' primer of the partially-randomized pool.
15 To allow for the possibility that the 5' primer formed part of the aptamer core, the original 5' primer sequence was included in the partially-randomized region of the new pool and a different 5' primer was appended for amplification. Because of differences in the relative
20 rates of phosphoramidite incorporation during DNA synthesis, a biased mix of all four nucleotides was prepared with molar ratios of 3:3:2:2 (A:C:G:T). This mix was added to pure phosphoramidite stocks (A and C: 64% pure stock, 36% random mix; G and T: 55% pure stock,
25 45% random mix) to yield mixed stocks for pool synthesis.

Twelve random bases were added to either end of this core sequence and new constant primers for PCR amplification were included. The synthesis of this 156 nucleotide DNA sequence yielded a pool containing 8×10^{13}
30 different molecules, which were transcribed to yield a pool of RNA molecules clustered in sequence space around the original biotin aptamer sequence. The total yield from the DNA synthesis was approximately 77 μg (1.52 nmole). The quality of the synthetic DNA was determined
35 by a primer extension assay, which showed that only 8.7%

- 38 -

of the DNA molecules could serve as full length templates for Taq polymerase. The pool thus contains $1.52 \times 10^{-9} \times 6.02 \times 10^{23} \times 0.087 = 8 \times 10^{13}$ distinct sequences.

This second RNA pool was used to identify
5 ribozymes able to enhance the rate of self-alkylation with the haloacetyl derivative, N-biotinoyl-N'-iodoacetyl-ethylenediamine (BIE; Molecular Probes, Eugene, OR). BIE is normally used to biotinylate proteins by reaction with free cysteine sulfhydryls. To
10 provide one potential internal substrate for the alkylation reaction, the doped pool was transcribed in the presence of excess 8-mercaptoguanosine, thus yielding RNAs containing a single free thiol in the 5'-terminal nucleotide. Following an overnight (15 hour) room
15 temperature incubation with 200 μ M BIE, RNAs that had undergone the self-biotinylation reaction were isolated by streptavidin agarose chromatography.

In particular, reaction with BIE was terminated by the addition of 100 mM β -mercaptoethanol, 5 mM EDTA, 0.3
20 M NaCl, 50 μ g tRNA (*E. coli*, RNase-free, Boehringer-Mannheim, Indianapolis, IN). After five minutes, the mixture was precipitated with 2.5 volumes ethanol on dry ice. After washing and resuspension, the RNA was applied to 0.5 ml of a 50% slurry of streptavidin agarose in wash
25 buffer (1 M NaCl, 10 mM NaHepes, pH 7.4, 5 mM EDTA) that had been washed with 50 μ g tRNA. After rocking 30 minutes to allow streptavidin-biotin binding, the mixture was transferred to a 10 ml-column and washed with 4 x 10 ml wash buffer and 2 x 10 ml distilled water.

30 RNA bound to streptavidin could be affinity eluted by first saturating the free biotin-binding sites with excess biotin and then heating in the presence of 10 mM biotin at 94°C for 8 minutes. Amplification of the resultant molecules (by reverse transcription, PCR, and
35 transcription) yielded a pool enriched for catalysts.

- 39 -

After three rounds of selection, an increase in the proportion of RNAs binding to the streptavidin was observed (FIG. 9B). By the fifth round, 10% of the RNA ligated the biotin substrate. To select for the most active catalysts, the incubation time was progressively shortened from 15 hours to 30 minutes to 1 minute. After eight rounds of selection, no further increase in activity was observed suggesting that the complexity of the starting pool had been exhausted. Sequencing individual clones from the selected pool showed that 50% of the ribozymes were very closely related and were derived from a single progenitor. One of these clones, BL8-6, catalyzes self-biotinylation at a rate of 0.001 min^{-1} in the presence of $200 \text{ } \mu\text{M}$ BIE.

The rate of self-biotinylation was determined by a time course experiment. ^{32}P -labelled RNA was first resuspended in incubation buffer (100 mM KCl, 10 mM Na-Hepes, pH 7.4, 5 mM MgCl_2) and allowed to equilibrate for 10 minutes at room temperature. $200 \text{ } \mu\text{M}$ BIE was added to the mixture and aliquots were subsequently removed after 0 to 120 minutes of incubation. Samples were quenched and affinity purified as described in Haugland, Molecular Probes Handbook of Fluorescent Probes and Research Chemicals. Aliquots were counted in a scintillation counter following ethanol precipitation (total RNA count) and following binding to streptavidin agarose (product RNA count); the ratio of these two counts is the fraction reacted.

Optimizing enzymatic activity: It seemed likely that the original RNA pool from which the BL8-6 ribozyme was derived might not saturate the space of biotin-ligating ribozymes. To test the possibility that appropriate additional mutations to the BL8-6 sequence might increase its catalytic activity, a third RNA pool was generated based on its sequence but with non-wild-type nucleotides

- 40 -

substituted at each position with 30% probability (FIG. 8D) (using methods described above). The selection for catalytic activity was repeated as described above, but with both the reaction incubation time and the BIE concentration progressively lowered to select for the most active enzymes. After eight rounds of selection (ending with a 1 minute incubation period at 10 μ M BIE), active clones from the pool were sequenced and assayed for catalytic activity. Ribozymes in this pool were uniformly more active than their BL8-6 progenitor, with one clone (BL2.8-7) catalyzing self-biotinylation at a rate of 0.05 min⁻¹ in the presence of 100 μ M BIE (one hundred fold more active than BL8-6).

Nature of the reaction product: The observation that BL8-6 ribozyme transcribed without 8-mercaptoguanosine catalyzed the self-biotinylation reaction as efficiently as the thiol-containing RNA indicated that some site other than the free thiol in the 8-mercaptoguanosine base at the 5'-end of the RNA might serve as nucleophiles for the alkylation reaction. However, the observation that BL8-6 ribozyme transcribed without 8-mercaptoguanosine catalyzed the self-biotinylation reaction as efficiently as the thiol-containing RNA indicated that some other site was being alkylated. To identify the reactive site, 5'-end labelled BL8-6 ribozyme that had reacted with BIE was subjected to alkaline hydrolysis, and the resultant ladder of molecules was affinity purified on streptavidin agarose. In particular, RNA was partially hydrolyzed by heating to 90°C for 7 minutes in the presence of 100 mM NaHCO₃, pH 9.0 and subsequently ethanol precipitated. After resuspending in wash buffer, biotin-labelled RNA was affinity purified as described by Haugland (*supra*). Purified non-biotinylated RNA was obtained from the initial flowthrough fraction from the streptavidin agarose slurry (prior to washing). Full length RNAs and

- 41 -

those with the approximately 60 3'-terminal nucleotides deleted were retained by the streptavidin whereas shorter molecules were not bound. This result maps the biotin attachment site to the region ...5'-⁹²GGACGUAAA¹⁰⁰-3'...

5 Alkylation at the N7 position of purines leads to RNA strand scission following treatment with sodium borohydride followed by aniline acetate (this reaction serves as the basis for the RNA chemical sequencing) (Peattie, Proc. Natl. Acad. Sci. USA 76:1760, 1979). RNA
10 incubated with BIE, purified on streptavidin-agarose, and treated in this manner was cleaved at G⁹⁶ (...GGACGUAAA...) (FIG. 10A). Briefly, RNA was dissolved in 1.0 M Tris-HCl, pH 8.2 and 0.2 M NaBH₄. Following a 30 min. incubation, the reaction was quenched with 0.6 M sodium
15 acetate/0.6 M acetic acid, pH 4.5, containing carrier tRNA. Following precipitation and rinsing, the RNA was treated with 1.0 M aniline/acetate, pH 4.5 at 60°C for 20 min. No G⁹⁶-specific cleavage was observed for RNA that had been exposed to BIE but not biotinylated (i.e. the
20 streptavidin flowthrough fraction). G⁹⁶ is therefore the alkylation site for the ribozyme.

To further characterize the alkylation product, the BL8-6 ribozyme was transcribed with [α -³²P]-GTP, thus labelling phosphates attached to the 5'-hydroxyl of all
25 guanosines in the RNA. Following reaction with BIE, biotinylated RNA was streptavidin-purified and subsequently digested to 5'-monophosphate nucleotides with snake venom phosphodiesterase I. Labelled RNA was diluted with 25 μ L 10 mM NaCl, 10 mM MgCl₂, 10 mM Tris-
30 Cl, pH 7.4, and 5 μ L phosphodiesterase I (Boehringer-Mannheim, Indianapolis, IN) and incubated for 20 hrs at 37°C. Thin layer ion exchange chromatography was carried out by spotting plates pre-run with water to remove excess salts and then developed with 6 M formic acid.
35 The PEI cellulose plates (J.T. Baker Co., Phillipsburg,

- 42 -

NJ) indicated the presence of a radioactive species in the streptavidin-purified RNA that was absent from the streptavidin-flowthrough RNA. This adduct migrated more rapidly than 5'-GMP in this TLC system, and co-migrated with 7-methyl GMP, suggesting that the adduct carries a positive charge, consistent with alkylation at N7 (FIG. 10A and FIG. 10B). Although the possibility of alkylation at N1 or N3 cannot be ruled out, alkylation at either of these sites would not be expected to lead to strand cleavage following aniline treatment, but would be expected to disrupt reverse transcription, thus preventing catalysts using these nucleophiles from being enriched during the *in vitro* selection procedure. Taken together, these results strongly suggest that N7 of G⁹⁶ is the alkylation site.

The catalyzed rate enhancement: The background rate of guanosine alkylation by BIE was determined by two independent methods. First, radiolabelled random sequence RNA (from the pool used to isolate the original biotin binder) was incubated for 24 hours with or without 200 μ M BIE. The specific increase in the fraction bound by streptavidin agarose (0.15%) after extensive washing was taken as a measure of the background reaction. Assuming an average of 28 guanosines/RNA sequence, this fraction corresponds to a non-catalyzed alkylation rate of $2.3 \times 10^{-6} \text{ s}^{-1} \text{ M}^{-1}$. In a similar approach, low concentrations of [α -³²P]-GTP were incubated overnight in the presence or absence of 200 μ M BIE and after 12 hours, affinity purified by streptavidin agarose. The fraction specifically bound (3.4×10^{-5}) indicates a non-catalyzed rate of $2.3 \times 10^{-6} \text{ s}^{-1} \text{ M}^{-1}$, in close agreement with that obtained from the RNA labelling experiment. A time course experiment with BL2.8-7 RNA yields a catalyzed biotinylation rate of approximately $8 \text{ s}^{-1} \text{ M}^{-1}$. The ribozyme rate enhancement is thus approximately 3×10^6 ,

- 43 -

comparable to that of the most active catalytic antibodies although substantially less than that of many natural protein enzymes (Tramontano et al., J. Am. Chem. Soc. 110:2282, 1988; Janda et al., ibid. 112:1275, 1990).

5 Structural differences between the biotin binder and the biotin ligator: Given that the biotin ligator arose by mutagenesis of the biotin binder sequence and that both molecules interact specifically with biotin, we expected to find significant structural similarities between the
10 two RNAs. Simple comparison of their primary sequences, however, failed to identify a well-conserved domain that might play a functional role; mutations appear randomly distributed along the length of the two sequences. To characterize the functional cores of the two molecules,
15 we analyzed the sequences of active clones isolated from the two mutagenized RNA pools generated from the biotin aptamer and self-alkylating ribozyme sequences. After four rounds of reselection with the biotin aptamer-derived pool, >40% of the applied RNA bound tightly to
20 biotin agarose. Similarly, three rounds of re-selection of the self-alkylating ribozyme-derived pool yielded a collection of RNAs with activity matching that of the original BL8-6 clone, and five additional rounds of selection increased the activity ~100-fold.
25 Approximately thirty individual RNAs from each of these subcloned pools were sequenced and analyzed to determine which nucleotide positions were conserved and which pairs of nucleotides covaried to maintain Watson-Crick base pairing. The results of these experiments are summarized
30 below and in FIG. 11, FIG. 12A, and FIG. 12B.

Two regions of the biotin binder are very highly conserved in clones that retain binding activity (FIG. 11). Mutations at the 5' and 3' ends of the first conserved domain (changing the A⁵³.G⁷⁰ pair to either C:G
35 or A:T) suggest a hairpin structure stabilized by a 4-

- 44 -

base-pair Watson-Crick duplex. Seven non-paired bases in the middle of the first domain directly complement the 3'-terminal half of the second conserved domain, thus suggesting a pseudoknot structure (FIG. 12A). In that

5 the bases in these conserved domains are essentially invariant, the sequence data provide no covariational evidence for the pseudoknot. To test the proposed structure, a series of site-directed mutants was generated and assayed for binding to biotin agarose.

10 Single-base substitutions that disrupt proposed Watson-Crick base pairs in the pseudoknot completely abolish biotin binding while compensatory second site mutations that introduce non-native Watson-Crick base pairs are able to largely restore biotin binding. These data

15 strongly support the proposed pseudoknot model for the biotin aptamer.

Comparison of the sequences of active ribozymes from the BL8-6 re-selection indicate a striking change in structure relative to the original biotin binder.

20 Nucleotides involved in the pseudoknot base-pairing (53-70, 101-107), virtually invariant in the biotin binders, are poorly conserved in the enzyme sequences (FIG. 11). In contrast, the ribozyme sequence in the region corresponding to the variable connecting loop of the

25 biotin binder (nucleotides 71 to 94) appears to be well conserved, suggesting a structural role. Nucleotides that are very highly conserved in the biotin binder but not involved in the pseudoknot base pairing (...5'-

95CGAAAAG¹⁰¹-3'...) are retained in the self-alkylating

30 enzymes but with a highly conserved change to ...5'-

95CGUAAAG¹⁰¹-3'... These results suggest that the change in function from biotin binding to alkylation of RNA with BIE is achieved by major structural rearrangements.

Further analysis of the BL8-6-derived sequences

35 suggested a cloverleaf structure with several remarkable

- 45 -

similarities to tRNA (FIG. 12B). The sequence ...5'-
94ACGUAAA¹⁰⁰-3'... is presented as the tRNA variable stem,
flanked on either side by extended duplexes (as indicated
by several observed Watson-Crick covariations). The
5 single guanosine in the variable stem serves as the
internal alkylation site for the enzyme. One
interpretation of these results is that the
hexanucleotide segments CGAAAA and CGUAAA directly
mediate the interaction with biotin in the biotin binder
10 and the biotin ligator respectively, although they are
presented in strikingly different secondary structure
contexts. Comparison of ribozyme sequences from the
third and eighth rounds of reselection suggest that the
increase in pool alkylation activity is achieved by
15 optimization of Watson-Crick base pairing in the
cloverleaf duplexes and an increased fraction of purines
(particularly adenosine) in the loop that caps helix 3.

To test the cloverleaf model for the biotin
ligator, a synthetic ribozyme was designed by modifying
20 one of the re-selected sequences such that 1) primer
sequences at the 5'- and 3'- ends not involved in the
cloverleaf were deleted; 2) non-conserved bulges in the
putative helices were removed, and 3) the variable loop
of approximately 45 nucleotides was replaced by a three
25 nucleotide loop sequence. The predicted lowest energy
structure for the resulting 99-nucleotide molecule is
shown in FIG. 13. This highly simplified structure has
~10 fold lower activity than the best re-selected clone,
but is still ~10 fold more active than the original BL8-6
30 ribozyme, thus supporting the proposed cloverleaf
structure (FIG. 13).

Two-component ribozyme: For a ribozyme to properly
qualify as an enzyme, it must emerge from the catalyzed
reaction unmodified. The self-alkylating ribozyme, which
35 has been selected to covalently modify its own active

- 46 -

site, fails to meet this requirement. The cloverleaf secondary structure, however, immediately indicates a way to engineer the ribozyme into two self-associating parts, one of which (BL-S) can function as a substrate for biotinylation while the other (BL-E) acts as a true enzyme (FIGS. 14A and 14B). A low level of BL-S biotinylation, corresponding to the non-catalyzed rate of alkylation was observed in the absence of BL-E. The initial rate of biotinylation of the RNA substrate increased linearly with increasing concentrations of BL-E, although the concentration of product never exceeded the concentration of enzyme. This result indicates that the two RNA pieces can associate with the BIE substrate to form a ternary complex capable of true catalysis. The extensive Watson-Crick base-pairing that drives complex formation most likely prevents dissociation of the biotinylated product and thus limits the enzyme fragment to a single catalytic event. Destabilizing the enzyme-substrate duplexes should make it possible to form a kinetically reversible complex that will dissociate after substrate biotinylation, allowing multiple rounds of turnover.

USE

Nucleic acids produced by the method of the invention can be used as *in vitro* or *in vivo* catalysts. In some cases the nucleic acids may be used to detect the presence of the ligand. For example, the nucleic acid may bind the ligand and catalyze a reaction which converts the ligand into a readily detectable molecule. The ribozymes created by the method of the invention can also be used in assays to detect molecules modified by the ribozymes which are not themselves ligands, e.g., an RNA phosphorylated by a polynucleotide kinase ribozyme.

- 47 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Szostak, Jack W.
Lorsch, Jon R.
Wilson, Charles
- (ii) TITLE OF INVENTION: NOVEL RIBOZYMES AND NOVEL RIBOZYME
SELECTION SYSTEMS
- (iii) NUMBER OF SEQUENCES: 91
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Fish & Richardson
 - (B) STREET: 225 Franklin Street
 - (C) CITY: Boston
 - (D) STATE: Massachusetts
 - (E) COUNTRY: U.S.A.
 - (F) ZIP: 02110-2804
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30B
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/299,498
 - (B) FILING DATE: 01-SEP-1994
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Clark, Paul T.
 - (B) REGISTRATION NUMBER: 30,162
 - (C) REFERENCE/DOCKET NUMBER: 00786/245001
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (617) 542-5070
 - (B) TELEFAX: (617) 542-8906
 - (C) TELEX: 200154

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 21 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGAACCUCUA GGUCAUUAAG A

21

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 18 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

- 48 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ACGUCAGAAG GAUCCAAG

18

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 110 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGAACACTAT CCGACTGGCA NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN

60

NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNCCTTGGTC ATTAGGATCG

110

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGUGGGAAGA AACUGCAGCU UCGGCUGGCA CC

32

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 134 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN CGAGGGAAGA AACUGCGGCA

60

CCNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNAGUGCCGG CUCGNNNNNN NNNNNNNNNN

120

NNNNNNNNNN NNNN

134

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 127 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

TGATTTCGCTA GCACGTCATT GGCTGGTAAC ACATGACACT ATACGAGCGA AAAA ACTACG

60

GCACCCTGGT CCGTTAGGGA CAACGACTAA AGTTAGTGCC CACGGGGCTC GTTCAGGGGG

120

GGCACGG

127

- 49 -

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 115 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

AGTCTCGCTA GCACCTTATT GGCTGGTAAC ACCTGACACT ATACGAGCGA AAAA	60
CTACTGCTGGT CCGTACGGGC CATGGACTTA AGATAGTGCC CACGGGGCTC GTTCA	115

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 127 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GGACTCACTA GCACGTTGTT GGCTGGTAAC ACCCGACCCT ATACGAGCGA AAAA	60
CTACTGCTGGT CCATACGGGA CTTGGACTAA AGTTAGTGCC CACGGGGCTC GTTCAGGGGG	120
GGCACGG	127

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 127 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

AGACTCACTA GCGCGTTATT GGCTGGTAGC CCCTGACACT ATACAGCGAA AATACTGCGG	60
CACCCTGGTC CGTACGGGAC ATGGACATTA TGTTAGTGCC CACGGGGCTC GTTCAGGGGG	120
GGCACGG	127

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 130 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGATATGTTG ATTCGCCCCA GCCTATAAAG TGACTCAATT CGAGGGACGC AACTACGGCA	60
CCGTCTATCT GAATCGGACG CGGAACCTGT GCCGTCTCTA CTCTAACGTT AGCGGAAAAC	120
GTGGGTTGCG	130

(2) INFORMATION FOR SEQ ID NO:11:

- 50 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 130 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

```
AGATGTGTCG ATTCGCCACA GCCAACAAAG CGGCCCAATT CGAGGGACGC AACTTCGGCA      60
CCGTCTATCA GAACGGGACG CGGTTCTAGT GCCGTCTCTA TCCTAACGTT AGCGGAAAAG      120
GAGGGTTGCG                                     130
```

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 130 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```
AGATGTGTTG ATTCGCCTCG GCCTGTTTAG TGACCAATTT CGAGGGACGC AACTTCGGCA      60
CCGTCTACCT GCAATAGACG AGGTACTTAT GCAGGCCCTA CTTTAACGTT AGCGGGAAAC      120
GAGGGTTGCG                                     130
```

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 123 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```
AGTCTACATG GAAGTTGTAC TATCTAAGTG TACTCACCAA AGACGAGGGC AGGAAATACG      60
GCACCATTTG CTACGCAAGG CCAAGTGCC CGGCGTCGTT TCAGAAAGGA TAACGTTAGC      120
CTG                                     123
```

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 123 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```
AGGCCACTTA GATGTCGCAC TATCTAAGCG TACACGCCAA TTACGAGGGC AGGAAATACG      60
GCACCTCCAG CTACGCAAGG CCCCAGTGCC CTGCCTCAGT TCGGAACGGA TAACGTTACC      120
CTG                                     123
```

(2) INFORMATION FOR SEQ ID NO:15:

- 51 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 121 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AGACCTCGTG TAAGTCGTAC TATCTAGGAG TGCACACGAA TACGAGGGCA GGAAATACGG 60
 CACCATAACT ACGCAAGGCC CAAGTGCCCG GCCTTGATTC AGAACGGATA ACGTTAGCCT 120
 G 121

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TTATTTTCGTT CGCACCCAGT GATCGCTCGG GACTGGGGCC TCCGCTAGGG AGGACATTGC 60
 GGCACCCAAA CGACCACACA GAACGTGCTA ACGATAGTGC CGGCTAGCAT CCGTGAATGA 120
 ACTGCTGCTG CTGGCG 136

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 135 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

AGAAGTTGTT CGCACCCAGT GAACGCTCGG GACTGGGGCC TCCGCTAGGG AGGACATTGG 60
 GCACCCGAAC TATCACTCAG AACGTGCTAT CGATATAGCC GGCTAGCACC TGATTATGAA 120
 CTGCTGCTGC TGGCG 135

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GGATATTGTT CGCACCTGCT GATCGCTTGG GACTGGGGCC TCCGCTAGGG AGGACATTGC 60
 GGCACCCAAA CTATCACTCA GAACGTGCTA ACGATAGTGC CGGCTAGCTT CTGTAAGTGA 120
 ACTGCTGCTG TTGGCG 136

(2) INFORMATION FOR SEQ ID NO:19:

- 52 -

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 137 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

```
AGACCTTAAT TCGAAAGCGT ATTCAACTTA CCATATCTCG CGCCGAGGGA AGGACCATCG      60
GCGCCAACTA CAGAGCCGTG GTTAGCGGAC TCCGCAGTGC CGGCTCGGGG AATAGGGTTC      120
TCACGAATTA CCGGCAT                                     137
```

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 137 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

```
AGGCCTTAAT TCGAAAGCGT ATTCGACATA CCATATTTTG CGCCGAGGGA AGATCCTTCG      60
GCACAGACTA CAGCGTCGAG GTGAGCGGCG CACACTGTGT CGGCTCGGGG AATAGGGTTC      120
TCACGAATTA CCGGCAT                                     137
```

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

```
AGATGTGGTT GCATAGTAGG CAGCCGGGCA CTTACGCCGA ATCGAGGGAC GAGACCGGAG      60
CACCACGATG CGCCGCGATA CCTCATTTGG GATTAGTGCC GGCTAGGAAA GTGAGTTCCT      120
TATGACCTGC CTCCAC                                     136
```

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

```
AGATGTGGCG GCATAGTAGG CAGCCGAGCA CTAACGCCAA ATCGAAGGAC GAGACTGCGG      60
CTCCACGATG CGCCGCGATG CCACTTTTGA GATTAGTACC GGCTGGGAAA GTGAATTCCT      120
TCTGGCCTGT CTCCAC                                     136
```

(2) INFORMATION FOR SEQ ID NO:23:

- 53 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 137 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

AGATCGATTG GAGACGCCCT GCGTACTTT AGGTAGAAAA CTCCGACGGA AAAAAGTGGC 60
GCACCGTGGG AGTAGAGGAT AGATAACAGG GCATTAGTGC CGGCCTCGCA AAGCTACCAT 120
GAGATGGAGC GATCAGG 137

(2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 137 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GGTTAGATTG GAAGCGCCCC GACTTACTTT AGGTTGAAAA CTCCGACGGA AAAACTACAG 60
CACCGTGGGA GTAGAGGATG GGATATCAGG CATTAGTGCC GGCCTCGTAA AGCTACCAGG 120
ATATTGGGAC GATCAGG 137

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 34 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

CGAGGGAAGA AAAUGCGGCA CCAGUGCCGG CUCG 34

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 41 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

ACGAGCGAAA AAACUACGGC ACUAGUGCCC ACGGGGCUCG U 41

(2) INFORMATION FOR SEQ ID NO:27:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

- 54 -

CGAGGGCAGG AAUACGGCA CCAGUGCCCG GCCUUG

36

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

GCUAGGGAGG ACAUUGCGGC ACCAGUGCCG GCUAGC

36

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 36 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CCGAGGGAAG AUCCUUCGGC ACAUGUGUCG GCUCGG

36

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 92 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

GGAACCUACG AGCGAAAAA CUACGGCACU CUGGUCCAUA CGGGACUUGG ACUAAAGUUA

60

GUGCCACGG GGCUCGUUCA AGGUUCUCAC GG

92

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 85 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ACGAGCGAAA AAACUACGGC ACUCUGGUCC AUACGGGACU UGGACUAAAG UUAGUGCCCA

60

CGGGGCUCGU UCAAGGUUCU CACGG

85

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 112 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

- 55 -

GGAACACTAT CCGACTGGCA CCNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN 60
 NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNCCTTGG TCATTAGGAT CG 112

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 156 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GGAGGCACCA CGGTCGGATC CNNNNNNNNN NNGGAACAC TATCCGACTG GCAAAGACCA 60
 TAGGCTCGGG TTGCCAGAGG TTCCACACTT TCATCGAAAA GCCTATGCTA GGCAATGACA 120
 TGGACTNNNN NNNNNNNNCC TTGGTCATTA GGATCG 156

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 156 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GGAGGCACCA CGGTCGGATC CGGTTTATTA TCATGAGCCC GACTCGACGG GCACTGTACA 60
 TAAGCTTCGG ATGCCATAGT TTAGACACTA TGGACGTAAA GCCCATGCTA GGCAAAGACA 120
 TTGACTGCAT GAGCGCCGCC TTGGTCATTA GGATCG 156

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

NNNNNNNNNN NNGGAACACT ATCCGACTGG CACCGACCAT AGGCTCGGGT TGCCAGAGGT 60
 TCCACACTTT CATCGAAAAG CCTATGCTAG GCAATGACAT GGACTNNNNN NNNNNNN 117

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 117 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

TTGCGGTGGG ANGGACCACA TGCCGCCTGG CACCGACCAT AGGCTCGGGT TGCCAGAGGT 60
 TCCACAGTTT CATCGAAAAG CCTATGCTAG GAGGTTACCT AGACTTAGGG GTTCACT 117

- 56 -

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

ATTTCGCGAT GGGGAGCACA TAGCAACTGG CACCGACCAT AGGCTCGGGT TGCAAGAGGT 60
TCCACACTTT CATCGAAAAG CCTATGCTAG GCAATGACAT GGACTNNNNN NNNNNNN 117

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

TCTTCGGAGG CCGTTACAGA CACACACTGG CACCGACCAT AGGCTCGGGT TGTGTGAGGT 60
TGCCCATGTT CATCGAAAAG CCTATGCTAC CCACTGACAT GGACTTTATC CACAAGT 117

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

CAGTTATTCT GCGTAACACA TTCTGACTGA CACCGACCAT AGGCTCGGGT TGCCCTAGTT 60
GCCACACTTT CAACGAAAAG CCTATGCTAA CCTATGACGT GGACTCCGGC ATGNNNN 117

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

CAAAGGTCCT ACGGAATACA CTCTAACTGA CACCGACCAT AGGCTCGGGT CTCCAAAGGT 60
GCCACATTTT CAGCGAAAAG CCTATGCTAT CCAATGGCAT GAAGTATCAC GTCTACT 117

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

- 57 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

NNCGCATCGT GCTGAAGACA TTCCGACTTC GACCGACCAT AGGCTCGGGT TCCCAAAGTT 60
GTCTCACATT CTTTGAAAAG CCTATGCTAC CTAGTGACAA GGATTACGCC CGCTGAG 117

(2) INFORMATION FOR SEQ ID NO:42:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

ACGTCCGCCA ACGGTGGACA TTCTGACGGG CACCGACCAT AGGCTCGGGT TGGCCGCGGT 60
TTCATACTTT CATTGAAAAG CCTATGCCAG GCAGTGACAT GAACTTTGAG GTAAAGT 117

(2) INFORMATION FOR SEQ ID NO:43:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

CCCTGTTAAA GAGGAACACA TTCCGACTGC TACCGACCAT AGGCTCGGGT TCGTTGAGGT 60
GCCACACATG CATTGACAAG CTTATGCTAG GGGTTGCCAT GGACTNNNNN NNNNNNN 117

(2) INFORMATION FOR SEQ ID NO:44:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

CAAGAACCGG CCGAAAAACA TTCCAATGG TACCGACCAT AGGCTCGGGT TCCAGACAT 60
TACACATTTT CTTTGAAAAG CCTATGATAT CCGCTGACCG TGACCGCTAG CGGCATC 117

(2) INFORMATION FOR SEQ ID NO:45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

TGGACTTTTC ACGGAACATG TTCCGATTGG CACCGACCAT AGGCTCGGCT TTCCAGAGGT 60
GCCACAACCTT CATTGAAAAG CCTATGCTAG CCAATGACCT GGACCATCAC AAAGGTT 117

(2) INFORMATION FOR SEQ ID NO:46:

- 58 -

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CTTCATTAAA GGGGAAAACA TTCCGACTGG GACCGACCAT AGGCTCGGTT TTTCAGAAGG 60
CACTCTGTTG CGTCGACAAG CCTATGCTGG ACCATGACCT GGACTATTTG CCCAGAT 117

(2) INFORMATION FOR SEQ ID NO:47:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TGATGAGAGC TACGAACACA CACCGACTGG CACCGACCAT AGGCTCGGTT TGCCTCAGAT 60
TCTTACCTTT CTTTGAAAAG CCTATGCTTG CTAATGACCT GGATTTGAGA ACANNNN 117

(2) INFORMATION FOR SEQ ID NO:48:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GACACAAAGC AGGCAACAAA TTCCGACTGG TACCGACCAT AGGCTCGGTT TGCCCCGAGCT 60
TCCACACTTT CATCGAAAAG CCTATGTTAG CTAATGACAG GGAGGACTCG ATGTGGT 117

(2) INFORMATION FOR SEQ ID NO:49:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

CCGAGCGGTC GGGGACGACA TTCCGTCTGG CACCGACCAT AGGCTCGGTT CTCCAGAGCT 60
TCCAAACCTT CTTGAAAAG CCTATGCTGG GCAATGACAT GGACTNNNNN NNNNNN 117

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

- 59 -

AGTGTCATAT TAGGGACACA GTCCGTATCG CACCGATCAT AGGCTCGGTT TGGCACGCGT 60
GCCACACTTG CAACGACAAG CCTATGGTAG TCCATAACCT GGACTACAAA CCCGATT 117

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

CCCTAGTGGG TAGGAACACA TTACGCCTGG CACCGACCAT AGGCTCGGTT GACCAGCGTT 60
TCCACACTTT CATCGAAAAG CCTATGCTTG CCATTGACAT GGACTCACGC ATTGCAT 117

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

GTGCCGACTT ACGGTTTACA TTCAAACCTGG CACCGACCAT AGGCTCGGTT TGCCTAACGT 60
TTCAAACCTTT CATCGAAAAG CCTATGCTTG GCAACGGTTA GGGTTTCGCA CGGCGAT 117

(2) INFORMATION FOR SEQ ID NO:53:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

CTGCACAGGT AGGGAACGCA TTTCGACTCG CACCGACCAT AGGCTCGGTC AGCGAGTTGC 60
GCCCCAATTT CAACGAAAAG CCTATGCTAG GTAATGCCAT GGACTGGTTC GTATCAT 117

(2) INFORMATION FOR SEQ ID NO:54:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GACGGAACCG TTTTAACACG TTCCGACCGG CACCGACCAT AGGCTCGGTT TGCCAGAGCT 60
TCACAACTTT CATCGAAAAG CCTATGAAAT GTAACGACAA GGACTACTCG ACCAGCA 117

(2) INFORMATION FOR SEQ ID NO:55:

- (i) SEQUENCE CHARACTERISTICS:

- 60 -

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

GGGTCGTGGC GCGGAACAAA TTCCACAGG CACCGACCAT AGGCTCGGTT TGCCTGTTGC 60
TCCACACCTT CATCGAAAAG CCTATGCCCC GCAATCACTT GGCCTTTGGA CGTCATT 117

(2) INFORMATION FOR SEQ ID NO:56:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

GCTCTGTTTCG GTTCAACAAA TTCACACTGG CAAAGACCAT AGGCTCGGTT TGCCAGAGGT 60
GCCACAGTTC ACTCGAAAAG CCTATGATCG CCAATGACAT GTACCTCACG CTAGGCA 117

(2) INFORMATION FOR SEQ ID NO:57:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

ACACTATGTA CTGGAAAACG TTCGGACACA CACCGACTAT AGGCTCGGTT TGCCATTGGT 60
GCCACAGTTG CAGCGAAAAG CCTATGCGGG GCCATGACAC GTACTGCCCA GTAACGT 117

(2) INFORMATION FOR SEQ ID NO:58:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:58:

TGCTACTGTT ATGTAACACA TTCCGACTGC GACCGACCAT AGGCTCGGTT TTCCAGACGT 60
TCGTCACTTG CTTGACAAG CCTATGAAAT TCAATGACAT GGCCTGGCTA GGCGCGA 117

(2) INFORMATION FOR SEQ ID NO:59:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:59:

- 61 -

TCTATGGCCG TGCAAACACA CTACGTCTGG CCCCACCACAT AGGCTCGGGT TGCCAGCGTT 60
TGCAAGGTTT CATCGAAAAG CCTATGCGAT CTAATGACAT GGACCGGAAG GCCCAAT 117

(2) INFORMATION FOR SEQ ID NO:60:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:60:

CTAAATTTGG TTGAAACACA TGCAGACTGG CCCCACCACAT AGGCTCGGGT TGTCAGAGGT 60
GCTTCACGTT CCTCGAAAAG CCTATGTGAT GGAATGACAT TGACTGAGGG ATGCGGT 117

(2) INFORMATION FOR SEQ ID NO:61:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:61:

GCNGAGGGCT CCGGTACACA TGCAGACTGG TCCCACCACAT AGGCTCGGGT TACCAGACCT 60
TCAACTACTT CTTGAAAAG CCTATGCCG TCAAGGCCAT GAACGCTCAA TCAGTGT 117

(2) INFORMATION FOR SEQ ID NO:62:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:62:

TGTCCGAACG ACGTATGCCA TTCCGTCTGG CCCCACCACAT AGGCTCGGAT TACCATTCTG 60
TACACACTTT CATCGAAAAG CCTATGCTGT TCAATGGCCC GGAATTCACT AGATGGT 117

(2) INFORMATION FOR SEQ ID NO:63:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:63:

CGGAATTACA CTGGATCACA TCCCAGCTGG CCCCACCACAT AGGCTCGGGT TGCCAGTGCT 60
TACACCTTT CACCGAAAAG GCTATGCTAG GCCATGCCAT TAACTNNNNN NNNNNN 117

(2) INFORMATION FOR SEQ ID NO:64:

- (i) SEQUENCE CHARACTERISTICS:

- 62 -

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:64:

GGTTTATTAT CATGAGCCCG ACTCGACGGG CACTGTACAT AAGCTTCGGA TGCCATAGTT 60
TAGACACTAT GGACGTAAAG CCCATGCTAG GCAAAGACAT TGA CTGCATG AGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:65:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:65:

GGTTTATCAT GTTTTAATCC CTACGCGGTC ACATTTGAAT AACCGGGGAA TTACAGAGTG 60
TAAACACTAT GAACGTAAAG ACCATGCGAA GCTATGACAC TGA CTGCATG GTCGCGG 117

(2) INFORMATION FOR SEQ ID NO:66:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:66:

GGGGTTTTTG TCGCGGACCC TCGCGACGTT CACTGTACAT AAGCTTCGGA TGCCGTAGAG 60
TAAACACTGC GGACGTAAAG CTCATGTTGG GTATTAAACC AAACAACATT AGCCCCG 117

(2) INFORMATION FOR SEQ ID NO:67:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:67:

AGCTTCTCAT CAGTCGGTCC CACTCCACCG ACATTTACGT AAGCTTTGGA TGCCATAGTA 60
AAAACACTAT GGACGTAAAG CGCAACGTAG CCCAAGATAT TGACAGTTTG AGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:68:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:68:

- 63 -

GTTTTATGTT CAAGTGCCCG AAACGGCCGG CACTGTACAT AACCTCGGA TGCAATAGTC 60
TAGACGCTAT TGGTGTAAG CCCATATTAG ACAAGGACCT TGTCTTCATG AGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:69:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:69:

GTTTATGAT TGTGAGCCCG GCTCCACGGT CACTCTGAAG ATGCTTCGGA TGCCATAGTT 60
CGCACACTAT GGACGTAAAG ATTGTTTCGAG TCACAGACAG TAGCTGCACA ATCGCCG 117

(2) INFORMATION FOR SEQ ID NO:70:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:70:

GGTTGAAATA AGCGTTAGGC CTACTTGACG CTCAGTAGGC AATCACCGBA TGCCGTAGTT 60
TATACACTAT GGACGTAAAG GTCATGCTGT TCTAAGACAT TGTCTGCATG ACCGCCG 117

(2) INFORMATION FOR SEQ ID NO:71:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:71:

GAAATTTTGT GTGCAGACAC TACTCTCCTG CACCGTTTAA AAGCTTCGGA TGCCATAGGT 60
TAAAAACTAT GGACGTAAAG CGCATGATCG GTAAACACAG TTACTGCATG ATCGCCG 117

(2) INFORMATION FOR SEQ ID NO:72:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:72:

GGTTTATCAT GTTTTAATCC CTACGCGGGT CACATTTGAA TACCGGGGAA TTACAGAGTG 60
TAAACACTAT GAACGTAAAG ACCATGCGAA GCTATGACAC TGACTGCATG GTCGCGG 117

(2) INFORMATION FOR SEQ ID NO:73:

- (i) SEQUENCE CHARACTERISTICS:

- 64 -

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:73:

GGTTTATCAT GTTTTAATCC CTACGCGGGT CACATTTGAA TACCGGGGAA TTACAGAGTG 60
TAAACACTAT GAACGTAAAG ACCATGCGAA GCTATGACAC TGA CTGCATG GTCGCGG 117

(2) INFORMATION FOR SEQ ID NO:74:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:74:

GGTTGAAAAA CATGAGCCAG TCTCGACGAG ACTTCTCGTT TCTAATCGGA TGCCATAGTT 60
AAGATACTAT GGACGTAAAG CGCTCGGTAG CTAAGAACAG TGTTTGCCAG CGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:75:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:75:

GGATTGTTAT ACCTTGGCCT GGATCCTAGC CACTGTAGCT ATCATCCGGA TGCCAGAGTT 60
TAGCCACTCT GGACGTAAAG CTCATGTTAA GAATAGACAT TGAATGCATG AGCGCCC 117

(2) INFORMATION FOR SEQ ID NO:76:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:76:

GATGCATTAT CTCGCGTGCG TGTAGACGGG GTCGACACGC AAGCTTCGGA TGCCATAGAT 60
TAGATACTAT GGACGTAAAG CTCATGTTAG TCAAAAACAC TGGCTCCATG AGCGCCC 117

(2) INFORMATION FOR SEQ ID NO:77:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:77:

- 65 -

GGAAAATCAT ATAAGTCCCG TCGCCCCGCG AACTTTACGT AAGATTCGGA TGCCATAGTT 60
TATCCACTAT GGGTGTAAG GTCATGCTAT ACCAACACAT TTATGGCATG ATCGCCG 117

(2) INFORMATION FOR SEQ ID NO:78:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:78:

GAAATTTTGT GTGCAGACAC TACCCTCCTG CACCGTTAAA AAGCTTCGGA TGCCATAGGT 60
TAAAAACTAT GGACGTAAAG CGCATGATCG GTAAACACAG TTACTGCATG TCGCCG 117

(2) INFORMATION FOR SEQ ID NO:79:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:79:

GGTGTATTAG CTTGAGTCCA ACTCCACGAG CACTATGAAT AATCTTCGGA TGCCATCGTT 60
TCAACACGAT GGACGTAAAG CCCACTGTTG GCAAATACAT TGAATGCAGG TCGCCG 117

(2) INFORMATION FOR SEQ ID NO:80:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:80:

GATGCATTAT CTCGCGTGCG TGTAACGCGG GATCGACACC AAGCTTCGGA TGCCATAGAT 60
TAGACACTAT GGACGTAAAG CGCATGTTAG TAGAAATCAA CTGCAGCACG ACCGCCG 117

(2) INFORMATION FOR SEQ ID NO:81:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 117 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:81:

GAAATTTTGT GTGCAGACAC TACTCTCCTG CACCGTTTAA AAGCTTCGGA TGCCATAGGT 60
TAAAAACTAT GGACGTAAAG CGCATGATCG GTAAACACAG TTACTGCATG TCGCCG 117

(2) INFORMATION FOR SEQ ID NO:82:

- (i) SEQUENCE CHARACTERISTICS:

- 66 -

- (A) LENGTH: 117 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:82:

GATTTATTCA TATGAGCCGG GTTGAAAGTA TAAAGTACTT TAGCTTCGGC TGCCAAAGTT 60
TATAAACTTT GGACGTAAAG CTCCTGCTTG GCAAATACAA AAGCTGCACG AGCGCCA 117

(2) INFORMATION FOR SEQ ID NO:83:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:83:

GGTTACTTAA TGCGACCAAC CTACGGGGCA CTGTCTACAT AAGTTTCGGA TGCCATAGTG 60
ATGCAACTAT GGACGTAAAG CCCATGCCAG ACTAAAACAT TGTCTGCATG CGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:84:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:84:

GGAGTCTTTT CATGAGTCCG ACTCTCCACT CATTGTTCAT AAGCTCCGGA TGCCATAGCT 60
CAAAAACATAT GGACGTAAAG CCCATGCTAA GCTCTCAAGT TGACTGCATG AGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:85:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:85:

GATTTATTCA TATGAGCCGG GTTGAAAGTA TAAAGTACTT TAGCTTCGGC TGCCAAAGTT 60
TATAAACTTT GGACGTAAAG CCCATGTTAG GTAAGATTAT TAACAGCATG TGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:86:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 117 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:86:

- 67 -

GCTTTATTCT CTCTTGCCCT GATCCACGGG CAGGATACGA GGGATGCGGA TGCCATATTT 60

TAAAAAGTAT GGACGTAAAG CCCATGATAA GCAAAGATTG TCACATCATG TGCGCCG 117

(2) INFORMATION FOR SEQ ID NO:87:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 99 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:87:

GGAACCAAGG CGGAUCCGGA UGAGAUCCGG AUGCCAUAGU AAAACACUA UGGACGUAAA 60

GCUCAGGCUG AAGACACAGC CUGAGCGCCG CCUUGGUUC 99

(2) INFORMATION FOR SEQ ID NO:88:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:88:

GGAACACUUAU GGACGUAAAG CUCAGGCUGA A 31

(2) INFORMATION FOR SEQ ID NO:89:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:89:

GGAAACAGCC UGAGCGCCGC CUUGGUUCGA AAGAACCAAG GCGGAUCCGG AUGAGAUCCG 60

GAUGCCAUAG UAA 73

(2) INFORMATION FOR SEQ ID NO:90:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 110 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:90:

GGAACACUUAU CCGAUGGCAC CGACCAUAGG CUCGGGUUGC CAGAGGUUCC ACACUUUCAU 60

CGAAAAGCCU AUGCUAGGCA AUGACAUGGA CUCCUUGGUC AUUAGGAUCG 110

(2) INFORMATION FOR SEQ ID NO:91:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 155 base pairs
- (B) TYPE: nucleic acid

- 68 -

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:91:

GGAGGCACCA CGGCUGGAUC CGGUUUAUUA UCAUGAGCCC GACUCGGGCA GCACUGUACA	60
UAAGCUCGGA UGCCAUAGUU UAGACACUAU GGACGUAAAG CCCAUGCUAG GCAAAGACAU	120
UGACUGCAUG AGCGCCGCCU UGGUCAUUAU GAUCG	155

- 69 -

CLAIMS

1. A method for producing a catalytic RNA molecule capable of binding a first ligand and catalyzing a chemical reaction modifying said RNA molecule,
5 comprising the steps of:

a) providing a first population of RNA molecules each having a first region of random sequence;

b) contacting said first population of RNA molecules with said first ligand;

10 c) isolating a first ligand-binding subpopulation of said first population of RNA molecules by partitioning RNA molecules in said first population which specifically bind said first ligand from those which do not;

15 d) amplifying said first ligand-binding subpopulation in vitro;

e) identifying a first ligand binding sequence;

f) preparing a second population of RNA molecules each of said RNA molecules comprising said
20 first ligand binding sequence and a second region of random sequence;

g) contacting said second population of RNA molecules with a second ligand capable of binding said first ligand binding sequence; and

25 h) isolating a subpopulation of said catalytic RNA molecules from said second population of RNA molecules by partitioning RNA molecules which have been modified in step g) from those which have not been modified.

30 2. The method of claim 1, wherein said first ligand is ATP or biotin.

3. The method of claim 1, wherein said second ligand serves as a substrate for said chemical reaction.

- 70 -

4. The method of claim 1, wherein said catalytic RNA molecule can transfer a phosphate from a nucleotide triphosphate to said catalytic RNA molecule.

5 5. The method of claim 4, wherein said transfer is to the 5'-hydroxyl or to an internal 2'-hydroxyl of said catalytic RNA molecule.

6. The method of claim 1, wherein said catalytic RNA molecule can transfer a phosphate from a nucleotide triphosphate to a nucleic acid other than said catalytic
10 RNA molecule.

7. The method of claim 6, wherein said nucleic acid is a ribonucleic acid.

8. The method of claim 1, wherein said first and second ligands are the same.

15 9. The method of claim 1, wherein said catalytic molecules can catalyze N-alkylation.

10. A catalytic RNA molecule which can transfer a phosphate from a nucleotide triphosphate to said catalytic RNA molecule.

20 11. The catalytic RNA molecule of claim 10, wherein said transfer is to the 5'-hydroxyl or to an internal 2'-hydroxyl of said catalytic RNA molecule.

12. A catalytic RNA molecule which can transfer a phosphate from a nucleotide triphosphate to a nucleic
25 acid other than said catalytic RNA molecule.

- 71 -

13. The catalytic RNA molecule of claim 12, wherein said nucleic acid is a ribonucleic acid.

14. A catalytic RNA capable of catalyzing N-alkylation.

5 15. A method for producing a catalytic RNA molecule capable of binding a first ligand and catalyzing a chemical reaction modifying a first substrate molecule bound to said catalytic RNA molecule, comprising the steps of:

10 a) providing a first population of RNA molecules each having a first region of random sequence;

 b) contacting said first population with said first ligand;

 c) isolating a first ligand-binding
15 subpopulation of said first population of RNA molecules by partitioning RNA molecules in said first population of RNA molecules which specifically bind said first ligand from those which do not;

 d) amplifying said first ligand binding
20 subpopulation in vitro;

 e) identifying a first ligand binding sequence;

 f) preparing a second population of RNA
molecules each of said RNA molecules comprising said first ligand binding sequence and a second region of
25 random sequence, each of said RNA molecules being bound to said first substrate molecule;

 g) contacting said second population of RNA molecules with a second ligand capable of binding said first ligand-binding sequence; and


30 h) isolating a subpopulation of said catalytic RNA molecules from said second population of RNA molecules by partitioning RNA molecules which are bound to a substrate molecule which has been modified in step

- 72 -

g) from those RNA molecules which are bound to a substrate molecule which has not been modified in step g).

16. The method of claim 15, wherein said second
5 ligand serves as a second substrate for said chemical reaction.

1/14



SEQ ID NO: 4

FIG. 1

2/14




FIG. 2

3/14




FIG. 3

4/14




FIG. 4

5/14

CLASS I

22 TGAATGCGCTAGCAGCTCATTTGGCTGGTGAACATGACACTATACAGCGGAAAACTACCGACCCCTGGTCCGTTAGGGACAAACGACTAAAGTTAGTCCACGGGGCTCGTTCAAGGTTCTCAGG
 45 AGCTGCGCTAGCAGCTCATTTAGGCTGGTGAACAGCTGACACTATACAGCGGAAAACTACCGACCCCTGGTCCGTTAGGGACAAACGACTAAAGTTAGTCCACGGGGCTCGTTCAAGGTTCTCAGG
 46 GGAATGCGCTAGCAGCTCATTTAGGCTGGTGAACAGCTGACACTATACAGCGGAAAACTACCGACCCCTGGTCCGTTAGGGACAAACGACTAAAGTTAGTCCACGGGGCTCGTTCAAGGTTCTCAGG
 57 AGACTACTAGCGCGCTTATTGGCTGGTGAACAGCTGACACTATACAGCGGAAAACTACCGACCCCTGGTCCGTTAGGGACAAACGACTAAAGTTAGTCCACGGGGCTCGTTCAAGGTTCTCAGG

SEQ ID NO: 6
 SEQ ID NO: 7
 SEQ ID NO: 8
 SEQ ID NO: 9

CLASS II

4' GGATATGTTGATTCGCGCCCGCCCTATTAAGTGACTCAATTTGAGGGACCGCACTACGGCACCCGCTTATCTGAAATCGGACCGGAACCTTGTCCTCTCTACTATACGTTAGCGGAAACGTTGCG
 25 AGATGCTGATTCGCGCACAGCCAAAGGCGCCCAATTTGAGGGACCGCACTACGGCACCCGCTTATCTGAAATCGGACCGGAACCTTGTCCTCTCTACTATACGTTAGCGGAAACGTTGCG
 43 AGATGCTGATTCGCGCCCTGCGCCCTGTTAGTGACCAATTTGAGGGACCGCACTACGGCACCCGCTTATCTGAAATCGGACCGGAACCTTGTCCTCTCTACTATACGTTAGCGGAAACGTTGCG

SEQ ID NO: 10
 SEQ ID NO: 11
 SEQ ID NO: 12

CLASS III

9 AGCTTACATGGAAGTTGTAATCTAAGTACTACCAAGACAGGCGAAGGAAATACGGACCTATTTGGTACGCAAGGCCCAAGTGGCCGGGCTTTCAGAAAGGATTAAGTTAGCTG
 16 AGGCCACTTGAATGTGCACTATCTAAGCGTACAGCGCAATTTACGAGGCGAGGAAATACGGACCTATTTGGTACGCAAGGCCCAAGTGGCCGGGCTTTCAGAAAGGATTAAGTTAGCTG
 42 AGACTCGCTTGAATGTGTAATCTAAGTACTACCAAGACAGGCGAAGGAAATACGGACCTATTTGGTACGCAAGGCCCAAGTGGCCGGGCTTTCAGAAAGGATTAAGTTAGCTG

SEQ ID NO: 13
 SEQ ID NO: 14
 SEQ ID NO: 15

CLASS IV

10 TTATTTGCTTGGCGCCAGTATGCTGCGACCTGGGCGCTCCCTGAGGGAGAACATTTGCCGACCAACCAAGCAGACAGAAAGTGGCTTAAGTCCGGCTAGCATCCGTGAATTAAGTCTGCTG
 14 AGAAGTTGCTTGGCGCCAGTATGCTGCGACCTGGGCGCTCCCTGAGGGAGAACATTTGCCGACCAACCAAGCAGACAGAAAGTGGCTTAAGTCCGGCTAGCATCCGTGAATTAAGTCTGCTG
 44 GAAATTTGCTTGGCGCCAGTATGCTGCGACCTGGGCGCTCCCTGAGGGAGAACATTTGCCGACCAACCAAGCAGACAGAAAGTGGCTTAAGTCCGGCTAGCATCCGTGAATTAAGTCTGCTG

SEQ ID NO: 16
 SEQ ID NO: 17
 SEQ ID NO: 18

CLASS V

1 AGACCTTAATTCGAAAGCGTATTCGACATACCAATATTTGGCGGAGGAAAGATCTCTGGCAGAGCTACAGAGTGGAGGTGAGCGGACACACTGTGTGTGGCTGGGAAATTAAGGTTCTCAGCAATTACCGGAT
 41 AGCCCTTAATTCGAAAGCGTATTCGACATACCAATATTTGGCGGAGGAAAGATCTCTGGCAGAGCTACAGAGTGGAGGTGAGCGGACACACTGTGTGTGGCTGGGAAATTAAGGTTCTCAGCAATTACCGGAT

SEQ ID NO: 19
 SEQ ID NO: 20

CLASS VI

13 AGATGCTGCTTGCATATGCTAGGACCGCCGGGCACTTACCGCAATTCGAGAGAGACCG - AGCACCACCATGCGCGCCGATATCTCATTTGGGATTAAGTCCGCTTAGGAAATTAAGCTTCTCTTAC
 49 AGATGCTGCGGATTAATGAGCGCCGAGACCTAACCCGAAATTCGAAAGAGAGCTGGGCTTCAGCATGCGCGCGATGCACTTTTGAATTAAGTACCTGCTGGGAAATTCGAAATTCCTTAATTCCTCTCTTAC

SEQ ID NO: 21
 SEQ ID NO: 22


CLASS VII

47 AGATGCTGCTGAGACCGCCCTGCGCTTCTTACGTAAGAAACCTCCGACCGGAAATTCGCGGACCTGGGAGTGAAGAGATTAAGTAAGG - CATTAAGTCCGGGCTCGCAAGCTACCATGAGATTAAGGCGATTAAG
 62 GGTAGATTTGGAAAGCGCCCGGACTTACTTAAAGTTGAAAACCTCGAGCGGAAA - CTACAGCAAGCTGGGAGTGAAGAGATTAAGTATACAGATTAAGTCCGGGCTCGTAAGCTACCATGAGATTAAGGCGATTAAG


SEQ ID NO: 23
 SEQ ID NO: 24

FIG. 5


ATP APTAMER CONSENSUS




CLASS I



CLASS III



CLASS IV



CLASS V






FIG. 6

SEQ ID NO: 30






SEQ ID NO: 31



Full-Length Cis-acting Ribozyme

Trans-acting Ribozyme



D

random N72 pool

SEQ ID NO: 32
 GGAACACTAT CCGACTGGCA CCNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN NNNNNNNNNN
 NNNNNNNNNN NNNNCTTGG TCATTAGGAT CG

blot in aptamer doped pool

SEQ ID NO: 33
 GGAGGCACCA CGGTGGGATC CNNNNNNNNN NNNNGAACAC taccgactg gcaaccgacca tagctcggg ttgccagag
 ttccacactt tcatcgaaaa gcctatgcta ggcactgaca tggactnnnn NNNNNNNNCG TTGGTCATTA GGATCG

blot in ligator doped pool

SEQ ID NO: 34
 GGAGGCACCA CGGTGGGATC Cggttatla tcatgagccc gactcgacgg gcaactglaca taagctcgg atgcatagt
 ttagacacta tgcagctaaa gcccatgcta gccaagaca ttgactgcat gagcgccgCG TTGGTCATTA GGATCG

9/14




FIG. 9A





FIG. 9B


FIG. 10A

NaBH₄ / aniline
workup

alkaline hydrolysis

10/14

FIG. 10B



blotting ligator clones

SEQ ID NO: 64
GGATTATTCATCAGCCCGACCTGCAGCCGGGACACTGTACACTTACGATTCGCATTAATTAGACACTATGGCCTAAAGCCCATGCTTAGGCCAAGAACAATTGACTGCATGAGCCGCC

SEQ ID NO: 65
-----C-GT-TTAAT-CCTA--CG-TCACA-T-GA---C-GGG-A-TA-G--G-A-----A-----G-A-T-T--C-----GT-G-

SEQ ID NO: 66
-GG-T--TGTCGG-A-CT-G-----TT-----G--AG-A-----GC-----T--G-T-TTA-ACCMA-AA-T--C--

SEQ ID NO: 67
A-C-C-C-C---GTC-GT-CCACTCCAGC-ACA-T--G-----AA-A-----G-ACG-C-C-T-----A-TT--

SEQ ID NO: 68
-T--GT--A-T-----AA--G-C-----C-C-----A-----G--T-GT-----ATG-T-G-T-C--G-AG-T--CA-T--

SEQ ID NO: 69
-T--GC-TAG--CG--C--T--C-GA-G-T-----CGC-----G--GTT-T-----G--G-T--C--G-T--

SEQ ID NO: 70
-G-AAAAGC-TTAGCCCTA-TTGA-C-T-A--GGC-T-AC-----G-----T--GT--GTT-T-----G--G-T--

SEQ ID NO: 71
-AAA-T--G-GTGC-A-ACTACTCT-CT--C--TT-A-----G--A-A-----G--A-C-T--C--G-T--

SEQ ID NO: 72
-G--T--T-A-GA-TGACT-AC-T--G-AG--A-G-GA-----AA-A-T-----G--A-C-T--C--G-T--

SEQ ID NO: 73
-----C-GT-TTAAT-CCTA--CG--T--AT-TG-AT-C-GGG-A-TA-G--G-A-----A-----G-A-T-T--C-----GT-G-

SEQ ID NO: 74
--G-AA-A-----A-T-----A-AC-T-C-GGT-CTTAA-----G-----A-T-----G-T-G--CT-GA--G-TT--CA-C--

SEQ ID NO: 75
-A-A-G-----ACCTG-G-TGAGTCTA-C-----GC-TCA-C-----G-----C-----T--T--A-A-T-----G-A-C--

SEQ ID NO: 76
-A-OC-----TC-C-TG--TG-A-----GT-QAC-GC-----A-----T--T--T--A--C-G-C--

SEQ ID NO: 77
-AAA-C-ATAAGT--T-G-CC-C-A--T--G--A-----TC-----GT--T--T--A--C-G-C--

SEQ ID NO: 78
-AAA-T--G-GTGC--A-CTAC-CT-CT--C-TA-A-----G--A-A-----G--A-C-T--C--G-T--

SEQ ID NO: 79
-----G-T--T-A-----C-A-----A-GA-----T-----CA--G--G-T--

SEQ ID NO: 80
-A-GC--TC-C-TG--TG-A-----G-TC-ACACG-----G-----G--G--T--

SEQ ID NO: 81
-AAA-T--G-GTGC--A-ACTACTCT-CT--C--TT-A-----G-----G--G--T--TAG-AT-ACTG-A--C-C--

SEQ ID NO: 82
-A-----CAT-----G-GT-GA-A-TAT-AA-----T-T-----A-----T-A-----T--T--MAG--C--A--T--C--

SEQ ID NO: 83
--AC--ATGC--C-AACCTA--GG-CA-TG-C-----T-----GAN-CA-----C-A-T-A-----A-TCTC-AG--

SEQ ID NO: 84
-AG-C-T-----T-----TC-ACT-T--T-----C-----C-C-A-A-----A-T-----A-T-----T--

SEQ ID NO: 85
-A-----CAT-----G-GT-GA-A-TAT-AA-----T-T-----C-----A-----T-A-----T-----T--GTT--A--T--

SEQ ID NO: 86
-C-----C-TCCT--TGA--C-----GGA--QAGG-A-G-----C-----A-----T-----A-A-----TTG-C--AT--T--

SEQ ID NO: 87




FIG. 13A




FIG. 13B





FIG. 14

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/10813**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C12P 19/34; C07H, 21/02

US CL : 435/91.31, 91.2; 536/23.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/6, 91.2, 91.3, 91.31, 91.2, 194, 195, 199; 536/23.1, 23.2; 514/44

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Trends in Biochemical Sciences, Volume 17, Number 3, issued 1992, Szostak "In Vitro Genetics", pages 89-93, see pages 91-92.	1-16
A, P	Biochemistry, Volume 34, Number 2, issued 1995, Huizenga et al, "A DNA Aptamer that Binds Adenosine and ATP", pages 656-665, see entire document.	1-15
A	Nature, Volume 346, issued 30 August 1990, Ellington et al, "In Vitro Selection of RNA Molecules that Bind Specific Ligands", pages 818-822, see entire document.	1-16

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search
07 NOVEMBER 1995

Date of mailing of the international search report

20 DEC 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

THOMAS G. LARSON, PH.D.

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/10813

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Biochemistry, Volume 31, Number 16, issued 28 April 1992, Pan et al, "In Vitro Selection of RNAs that undergo Autocatalytic Cleavage with Pb ²⁺ ", pages 3887-3895, see entire document.	1-16
X, P --- Y, P	Nature, Volume 374, issued 27 April 1995, Wilson et al, "In Vitro Evolution of a Self-Alkylating Ribozyme", pages 777-782, see entire document.	1-3, 9, 14-16 ----- 4-8, 10-13
Y	Science, Volume 261, issued 10 September 1993, Bartel et al, "Isolation of New Ribozymes from a Large Pool of Random Sequences", pages 1411-1418, see entire document.	1-8, 10-13, 15-16
Y	Science, Volume 257, issued 31 July 1992, Beaudry et al, "Directed Evolution of an RNA Enzyme", pages 635-641, see entire document.	1-16
Y	Nature, Volume 364, issued 05 August 1993, Sassanfar et al "An RNA Motif that binds ATP", pages 550-553, see entire document.	1-2, 4-7 10-13, 15
Y	Science, Volume 236, issued 19 June 1987, Cech, "The Chemistry of Self-Splicing RNA and RNA Enzymes", pages 1532-1539, see entire document.	1-8, 10-13, 15-16
Y	US, A, 5,093,246 (CECH ET AL) 03 March 1992, see entire document.	1-8, 10-13, 15-16
Y	US, A, 4,987,071 (CECH ET AL) 22 January 1991, see entire document.	1-8, 10-13, 15-16

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/10813

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/10813

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, BIOSIS, MEDLINE, CJACS (STN), Derwent World Patent Index.

Search Terms: catalytic RNA, ribozyme, polymerase chain reaction, PCR, selection, enrichment, enrich, kinase, phosphate transfer, ligase, hydrolase, alkylation, aptamere, ATP binding, biotin binding, Szostak Jack W, Lorsch Jon R, Wilson Charles.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-14, drawn to a catalytic RNA molecule and a first method of making said catalytic RNA molecule.

Group II, claims 15-16, drawn to a second method of making a catalytic RNA molecule.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The method for determining unity of invention under PCT Rule 13 permits, in addition to an independent claim for a product, an independent claim for a process of making said product and an independent claim for the use of said product. The first process of making is based on independent claim 1 and the second process of making is based on independent claim 15. Unity of invention exists between the product claims (10-14) and either the first process of making (claims 1-9) or the second process of making (claims 15-16), because the product and a process of making are considered to be linked by a "special technical feature." However, two processes for making the same product are not considered to be linked by a "special technical feature," therefore unity of invention does not exist between the first and second processes for making the claimed compound. Therefore, the claims are not so linked by a "special technical feature" within the meaning of PCT Rule 13.2 to form a single inventive concept.